



Section 4.55(1A) Modification Application Supporting Report

DA 24/964 – Kosciuszko Flow Trail Realignment,
Little Beauty Jumps Bypass and Wombat Walk
Connector Mountain Bike Trails

MOD 1 – Wombat Walk Connector Trail
Modification

Thredbo Alpine Resort,
Kosciuszko National Park

January 2025

Document Control

Revision	Date	Revision Type	Author	Approved by
A	20/11/2024	Draft	J.Best; C.Chalk	K.Delpit
0	20/12/2024	Final	J.Best; C.Chalk	K.Delpit
1	07/01/2025	Assessment of Section 4.55(1A) and 4.55(3) of the EP&A Act	J.Best	C.Chalk

Project: 24026MO
Kosciuszko Thredbo Pty Ltd

Contents

1	Introduction	1
2	Project Description.....	1
2.1	Approved Development under DA 24/964.....	1
2.2	Purpose of the MOD.....	1
2.3	Development components subject to this MOD.....	1
2.3.1	Construction Works.....	1
2.3.2	Condition Amendments	5
3	Statutory Framework	7
3.1	Environmental Planning and Assessment Act 1979	7
3.2	Section 100 of EP&A Act – Content of a Modification Application	8
4	Key Environmental Considerations.....	10
4.1	Biodiversity	10
4.2	Soil and Water	10
4.3	Aboriginal Cultural Heritage	10
5	Conclusion.....	12
6	References	12
7	Appendices.....	12

Figures

Figure 1: Site Plan.....	2
Figure 2: Photo Points.....	3
Figure 3: Waterfront land review (Source: NSW Government 2024).....	10

Tables

Table 1: Application Details	1
Table 2: Site Photo Points and Trail Description.....	4
Table 3: Superseded documents to be removed from Condition A.2	5
Table 4: Additional plans to be included in Condition A.2.....	6
Table 5: EP&A Act, Section 4.55 considerations	7
Table 6: EP&A Act, Section 4.15(1) considerations.....	8
Table 7: Content of modification application	9
Table 8: Aboriginal Cultural Heritage Due Diligence Assessment.....	11

1 Introduction

This Modification Application (MOD) relates to Development Application (DA) 24/964 which was granted development consent on 8 April 2024. This application is being made under section 4.55(1A) of the Act as the proposed changes to the Development are of minimal environmental impact.

Application details are provided in **Table 1**.

Table 1: Application Details

Details	
Applicant Name	Kosciuszko Thredbo Pty Ltd
Applicant Address	1 Friday Drive, Thredbo NSW 2625
ABN	ABN 95 000 139 015
Site address	2 Friday Drive, Thredbo NSW 2625
Land	Kosciuszko Flow Trail and Wombat Walk access track, Thredbo Alpine Resort, Kosciuszko National Park
Lot/Plan	876/DP1243112 (Thredbo Head Lease)
Consent Authority	Department of Planning, Housing and Infrastructure
Integrated bodies	The original development application was integrated development requiring assessment under the <i>Water Management Act 2000</i> in relation to the Little Beauty Jumps Park Bypass. The trail subject to this modification application is not located within waterfront land and therefore does not require assessment under the Act.
Development	Vegetation clearing Construction of mountain bike trail Site rehabilitation

2 Project Description

2.1 Approved Development under DA 24/964

As per the Development Consent for DA 24/964, the approved development was for construction of a Mountain Bike Trail – Upper N4 as outlined in Condition A.2.

2.2 Purpose of the MOD

Following completion of the Alpine Coaster installation it was determined the safety clearances for the original trail sections that traverse below the coaster infrastructure are not suitable. The purpose of the modification is to realign the trail away from the coaster footprint.


2.3 Development components subject to this MOD

2.3.1 Construction Works

Works subject to this MOD include (refer **Figure 1**) include:

- vegetation clearing
- construction of mountain bike trail
- site rehabilitation.

A description of the trail with photo point references is provided in **Figure 2** and **Table 2**.



Legend

- DA24/964 Proposed MOD
- - - DA24/964 Approved - Not built
- Wombat Walk - Existing
- Home Run
- Village Loop
- - - Close & Rehab
- 20m Construction Corridor

Scale: 1:1,378

9.750 9.5 19 28.5 38
 Meters

Map Projection: Universal
 Transverse Mercator
 Horizontal Datum: GDA 1994
 Grid: GDA 1994 MGA Zone 55




SITE PLAN

Project: Wombat Walk
 Connector DA24/964 MOD 1

Revision: A

Date: 9/12/2024

Produced By: BB



Legend

- DA24/964 Proposed MOD
- Wombat Walk - Existing
- Home Run
- Village Loop
- Close & Rehab
- Photo Points

Scale: 1:1,378

9.750 9.5 19 28.5 38

Meters

Map Projection: Universal Transverse Mercator
Horizontal Datum: GDA 1994
Grid: GDA 1994 MGA Zone 55

N

PHOTO POINTS

Project: Wombat Walk
Connector DA24/964 MOD 1

Revision: A

Date: 9/12/2024

Produced By: BB

Table 2: Site Photo Points and Trail Description




Photo Point ID / Description	Photo
PH1: The trail continues off the existing Wombat Walk access track onto the disturbed ski slope towards the summer access road.	
PH2: The trail traverses the edge of the disturbed ski slope	
PH3: The trail traverses the disturbed ski slope uphill of the tree island before connecting onto the existing Village Loop trail	

Photo Point ID / Description	Photo
PH4: The lower section of trail starts off the existing Home Run trail	
PH5: The trail traverses across the open ski slope, crossing below the tree island onto Sundance ski run before turning back towards Valley Terminal.	

2.3.2 Condition Amendments

2.3.2.1 Remove superseded documents in Condition A.2

The Development no longer requires the installation of a platform on Wombat Walk Connector. The Geotechnical Assessment and Form 4 was prepared subject to the installation of the platform. Trail construction works involved minor earthworks not involving excavations or fill in excess of one metre in vertical height. Therefore, geotechnical assessment of the trail is not required.

KT suggests removing the documents listed in **Table 3** from Condition A.2 in the modified consent.

Table 3: Superseded documents to be removed from Condition A.2

Ref No.	Document	Title/ Description	Author/ Prepared by	Date	Document Reference
2	Report	Site Environmental Management Plan (SEMP) Kosciuszko Flow Trail Realignment, Little Beauty Jumps Bypass and Wombat Walk Connector Mountain Bike Trails	Kosciuszko Thredbo Pty Ltd	5 March 2024	Rev 1

Ref No.	Document	Title/ Description	Author/ Prepared by	Date	Document Reference
7	Plan	Site Plan Wombat Walk Connector	Kosciuszko Thredbo Pty Ltd	16 October 2023	Rev 3
8	Plan	Typical Modular Bridge	Camstruct Consulting Pty Ltd	14 November 2023	23045-S01 A
9	Plan	Typical Modular Bridge Handrail	Camstruct Consulting Pty Ltd	14 November 2023	23045-S02 1
10	Geotechnical Assessment	Proposed Wombat Walk Connector Thredbo, NSW	ACT Geotechnical Engineers Pty Ltd	November 2023	AS/C14767. 2
11	Geotechnical Policy Form 4	Geotechnical Policy – Kosciuszko Alpine Resorts Form 4 – Minimal Impact Certification	ACT Geotechnical Engineers Pty Ltd	5 December 2023	-

2.3.2.2 Include additional documents in Condition A.2

This application is supported by the documentation listed in **Table 4**.

Table 4: Additional plans to be included in Condition A.2

Document	Title/ Description	Author/ Prepared by	Date	Document Reference
Plan	Site Plan Wombat Walk Connector DA 24/964 MOD 1	Kosciuszko Thredbo Pty Ltd, BB	9/12/2024	Rev A
Report	Wombat Walk Connector Realignment – Thredbo Alpine Resort – Flora and Fauna Assessment	Eco Logical Australia	20/12/2024	Version 1
Report	Site Environmental Management Plan Kosciuszko Flow Trail Realignment, Little Beauty Jumps Bypass and Wombat Walk Connector Mountain Bike Trails	Kosciuszko Thredbo Pty Ltd	20/12/2024	Rev 2
Report	Flow Trail Realignment Little Beauty Jumps Park Bypass & Wombat Walk Connector DA 24/964 MOD 1 Detailed Rehabilitation and Monitoring Plan	Kosciuszko Thredbo Pty Ltd	20/12/2024	Rev 0

2.3.2.3 Remove Part B requirements

KT suggests removing Part B requiring a construction certificate from the modified consent as the Development no longer includes the installation of a platform.

3 Statutory Framework

The Statement of Environmental Effects –Flow Trail Realignment, Little Beauty Jumps Pass to Wombat Walk Connector (KT 2024) provided a review and assessment against key legislation and planning instruments applicable to the Development. The Development remains substantially the same as the approved development under DA 24/964, including modification of the location of the mountain bike trail. Additional statutory considerations relevant to this MOD are provided below.

3.1 Environmental Planning and Assessment Act 1979

This MOD is being made under Section 4.55(1A) of the *Environmental Planning and Assessment Act 1979* (EP&A Act). A review of the Development against the relevant provisions in Section 4.55 is provided in **Table 5**.

Table 5: EP&A Act, Section 4.55 considerations

EP&A Act, Section 4.55	Comment
(1A) Modifications involving minimal environmental impact A consent authority may, on application being made by the applicant or any other person entitled to act on a consent granted by the consent authority and subject to and in accordance with the regulations, modify the consent if—	
(a) it is satisfied that the proposed modification is of minimal environmental impact, and	The MOD involves minimal environmental impact.
(b) it is satisfied that the development to which the consent as modified relates is substantially the same development as the development for which the consent was originally granted and before that consent as originally granted was modified (if at all), and	The consent as modified by this application is substantially the same development for which consent was originally granted.
(c) it has notified the application in accordance with— (i) the regulations, if the regulations so require, or (ii) a development control plan, if the consent authority is a council that has made a development control plan that requires the notification or advertising of applications for modification of a development consent, and	The application will be notified in accordance with the regulations.
(d) it has considered any submissions made concerning the proposed modification within any period prescribed by the regulations or provided by the development control plan, as the case may be.	Consideration will be given to submissions made.
Subsections (1), (2) and (5) do not apply to such a modification.	-
(3) In determining an application for modification of a consent under this section, the consent authority must take into consideration such of the matters referred to in section 4.15(1) as are of relevance to the development the subject of the application. The consent authority must also take into consideration the reasons given by the consent authority for the grant of the consent that is sought to be modified.	Refer Table 6 .

A review of the Development against the provisions in Section 4.15(1) of the EP&A Act is provided in **Table 6**.

Table 6: EP&A Act, Section 4.15(1) considerations

Environmental Planning and Assessment Act 1979, Section 4.15(1)		Comment
(1) Matters for consideration—general In determining a development application, a consent authority is to take into consideration such of the following matters as are of relevance to the development the subject of the development application— (a) the provisions of—		
(i) any environmental planning instrument		The Precincts – Regional SEPP is the only environmental planning instrument which applies to the site for this proposal.
(ii) any proposed instrument that is or has been the subject of public consultation under this Act and that has been notified to the consent authority (unless the Planning Secretary has notified the consent authority that the making of the proposed instrument has been deferred indefinitely or has not been approved)		Not applicable. There are no draft Environmental Planning Instruments that are applicable to the Development.
(iii) any development control plan		Not applicable. There are currently no development control plans applicable to the site.
(iia) any planning agreement that has been entered into under section 7.4, or any draft planning agreement that a developer has offered to enter into under section 7.4		Not applicable. There are no planning agreements applicable to Thredbo under the Precincts – Regional SEPP.
(iv) the regulations (to the extent that they prescribe matters for the purposes of this paragraph)		The DA and supporting information have been prepared in accordance with the relevant requirements of the EP&A Regulation.
the likely impacts of that development, including environmental impacts on both the natural and built environments, and social and economic impacts in the locality		The Development will result in acceptable environmental impacts, refer Section 2 .
the suitability of the site for the development		The site is considered suitable for the Development as it is located within predominately disturbed land.
any submissions made in accordance with this Act or the regulations		Consideration will be given to submissions made.
the public interest.		The trail modification is considered within the public interest as it will connect riders from the western side of the trail network to the bottom of the lifts at Valley Terminal. With the planned upgrade of Snowgums Chairlift to a new 6-seat MTB compatible lift, a linking trail from the western side of the trail network will allow riders direct access to the Snowgums bottom station. This will help to alleviate the rider traffic and congestion often experienced at the bottom of the Kosciuszko Express chairlift during peak visitation periods.

3.2 Section 100 of EP&A Act – Content of a Modification Application

A cross-reference to the requirements for a modification application in accordance with Section 100 of the *Environmental Planning and Assessment Regulation 2021* (EP&A Regulation) is provided in **Table 7**.

Table 7: Content of modification application

Content of modification application (Section 100 of EP&A Regulation)	Response
(1) A modification application must contain the following information—	
(a) the name and address of the applicant,	Kosciuszko Thredbo Pty Ltd 1 Friday Drive, Thredbo NSW 2625
(b) a description of the development that will be carried out under the development consent	A description of the development is provided in Section 2.3 .
(c) the address and folio identifier of the land on which the development will be carried out,	Refer Table 1
(d) a description of the modification to the development consent, including the name, number and date of plans that have changed, to enable the consent authority to compare the development with the development originally approved,	A description of the modification is provided in Section 2.3 . Documents to be superseded and additional documents are outlined in Section 2.3.2 .
(e) whether the modification is intended to— (i) merely correct a minor error, misdescription or miscalculation, or (ii) have another effect specified in the modification application,	Refer Section 2.3 . The Development will remain substantially the same as the development originally approved.
(f) a description of the expected impacts of the modification,	Refer Section 2 .
(g) an undertaking that the modified development will remain substantially the same as the development originally approved,	See comment against (e).
(h) for a modification application that is accompanied by a biodiversity development assessment report—the biodiversity credits information,	Not applicable, no BDAR prepared for original DA.
(i) if the applicant is not the owner of the land—a statement that the owner consents to the making of the modification application,	Owners Consent has been provided separately as part of this MOD.
(j) whether the modification application is being made to— (i) the Court under the Act, section 4.55, or (ii) the consent authority under the Act, section 4.56.	This MOD is being made to the consent authority under the EP&A Act. This MOD is substantially the same as the development for which the consent was originally granted, and results in acceptable environmental impact.
(2) Subsection (1)(i) does not apply if the consent of the owner is not required under section 98.	Not applicable
(3) If a modification application under the Act, section 4.55(1A) or (2) relates to BASIX development, or BASIX optional development if the development application was accompanied by a BASIX certificate, the application must be accompanied by—	Not applicable

4 Key Environmental Considerations

The impacts of the trail modification are consistent with those identified in the original Statement of Environmental Effects prepared by Kosciuszko Thredbo Pty Ltd (2024). With the implementation of appropriate environmental controls and sustainable trail building techniques the impacts of the modified trail are considered acceptable.

4.1 Biodiversity

A Flora and Fauna Assessment has been prepared by Eco Logical Australia to support this application. A copy of the report is provided in **Appendix A**.

4.2 Soil and Water

The Development will be constructed to effectively manage erosion and run-off in accordance with sustainable trail design concepts and construction techniques such as rolling contours, outslope, the half rule and 10% average guideline and use of frequent grade reversals to minimise erosion and soil stability risks. Where areas of disturbance do not form part of the final trail alignment, they will be stabilised and/or revegetated in accordance with the rehabilitation and monitoring which will assist in achieving an erosion resistant state.

The new section of trail is not located within 40 m of a watercourse (waterfront land), refer **Figure 3**. No further assessment is required.




Figure 3: Waterfront land review (Source: NSW Government 2024)

4.3 Aboriginal Cultural Heritage

An assessment of the modified trail section against the *Due Diligence Code of Practice for the Protection of Aboriginal Objects in New South Wales* (DECCW 2010) is provided in **Table 8**.

Table 8: Aboriginal Cultural Heritage Due Diligence Assessment

Step in Due Diligence Process	Comment
1) Will the activity disturb the ground surface or any culturally modified trees?	The Development will result in ground disturbance.
2) Are there any: <ul style="list-style-type: none"> a. relevant confirmed site records or other associated landscape feature information on AHIMS? And/or b. Any other sources of information of which a person is already aware? And/or c. landscape features that are likely to indicate presence of Aboriginal objects? 	<p>A search of the Aboriginal Heritage Information Management System (AHIMS) was undertaken on 20 Nov 2024. The search results (refer to appendices) identified no Aboriginal sites are recorded in or near site or surrounds. The site consists of steep terrain descending down the upper/mid slopes.</p> <p>Past Traces prepared an assessment for the proposed trail in 2023. The assessment concluded the works could proceed without further assessment as no Aboriginal or historical heritage sites (objects or places) have been identified within the project area. The potential for impacting on unrecorded heritage sites within the project area was assessed as extremely low, based on landform analysis and field survey (Past Traces 2023).</p> <p>Several historical independent assessments have been undertaken within the resort by Past Traces Heritage Consultants (2017-2023), NGH Environmental (2017), Iron Bark (2013), and URS Australia Pty Ltd (2004; 2005), as well as other due diligence assessments undertaken for original Development applications in the area. All studies concluded that the ski slope areas hold low potential for Aboriginal heritage sites. The studies also concluded that given the steepness and exposed aspect/lack of sheltering tors, the ski slopes are unlikely to have been favourable campsite locations.</p> <p>No landscape features that are likely to indicate presence of Aboriginal objects were identified within the site. It is considered the Development has low potential to impact on unrecorded Aboriginal objects or sites. There is no requirement to move onto Steps 3 and 4.</p>
3) Can harm to Aboriginal objects listed on AHIMS or identified by other sources of information and/or can the carrying out of the activity at the relevant landscape features be avoided?	Not applicable
4) Does a desktop assessment and visual inspection confirm that there are Aboriginal objects or that they are likely?	Not applicable

As identified above, all reasonable steps have been undertaken to ensure the Development fulfils the requirements of the Aboriginal Cultural Heritage Due Diligence Process. Potential impacts from the Development on objects or sites of Aboriginal Cultural Heritage significance are considered unlikely. In the unlikely event that Aboriginal objects are discovered, management measures outlined in the SEMP will be implemented.

5 Conclusion

This application has considered the potential impacts of the works in accordance with relevant statutory requirements. The modification to the original development will result in acceptable environmental impact, where the development will remain substantially the same as the development that was originally approved under DA 24/964.

6 References

Eco Logical Australia 2024, Wombat Walk Connector Realignment – Thredbo Alpine Resort – Flora and Fauna Assessment. Prepared for Kosciuszko Thredbo Pty Ltd.

Kosciuszko Thredbo Pty Ltd, 2021. Statement of Environmental Effects –Flow Trail Realignment, Little Beauty Jumps Pass to Wombat Walk Connector.

Past Traces, 2023. Aboriginal Cultural Heritage Due Diligence Assessment. Kosciuszko Flow Trail Realignment and Wombat Walk Connector.

7 Appendices

Appendix A – Flora and Fauna Assessment

A stylized topographic map with green contour lines is positioned on the left side of the page, extending from the top left towards the bottom left.

Wombat Walk Connector Realignment – Thredbo Alpine Resort – Flora and Fauna Assessment

Kosciuszko Thredbo Pty Ltd

DOCUMENT TRACKING

Project Name	Wombat Walk Connector Realignment – Thredbo Alpine Resort – Flora and Fauna Assessment
Project Number	24NAR9738
Project Manager	Ryan Smithers
Prepared by	Ryan Smithers
Reviewed by	Dave Coombes
Approved by	Ryan Smithers
Status	Final
Version Number	1
Last saved on	20 December 2024

This report should be cited as ‘Eco Logical Australia 2024. *Wombat Walk Connector Realignment – Thredbo Alpine Resort – Flora and Fauna Assessment*. Prepared for Kosciuszko Thredbo Pty Ltd.’

ACKNOWLEDGEMENTS

This document has been prepared by Eco Logical Australia Pty Ltd with support from Kosciuszko Thredbo Pty Ltd

Disclaimer
This document may only be used for the purpose for which it was commissioned and in accordance with the contract between Eco Logical Australia Pty Ltd and Kosciuszko Thredbo Pty Ltd. The scope of services was defined in consultation with Kosciuszko Thredbo Pty Ltd, by time and budgetary constraints imposed by the client, and the availability of reports and other data on the subject area. Changes to available information, legislation and schedules are made on an ongoing basis and readers should obtain up to date information. Eco Logical Australia Pty Ltd accepts no liability or responsibility whatsoever for or in respect of any use of or reliance upon this report and its supporting material by any third party. Information provided is not intended to be a substitute for site specific assessment or legal advice in relation to any matter. Unauthorised use of this report in any form is prohibited.

Contents

1. Introduction	1
1.1 The proposal	1
1.2 Direct and indirect impacts	2
1.3 Subject site, study area and locality	2
1.4 Biodiversity Offset Scheme	9
1.5 Topography, geology and soils	9
1.6 Disturbances	9
1.7 Planning and legislation	9
1.7.1 Environmental Planning and Assessment Act 1979	9
1.7.2 Biodiversity Conservation Act 2016	9
1.7.3 State Environmental Planning Policy (Precincts—Regional) 2021	10
1.7.4 Environment Protection and Biodiversity Conservation Act 1999	10
2. Methods	11
2.1 Database and literature review	11
2.2 Field surveys	11
2.2.1 Flora surveys	11
2.2.2 Fauna surveys	11
3. Results	13
3.1 Database and literature review	13
3.2 Flora	13
3.2.1 Subalpine Woodland	13
3.2.2 Exotic Grassland and Disturbed Ski Slopes	13
3.3 Fauna	15
3.3.1 Fauna habitats	15
4. Impact assessment	16
4.1 Impacts on vegetation communities	16
4.1.1 Subalpine Woodland	16
4.2 Impacts on threatened ecological communities	16
4.3 Impacts on flora species of conservation significance	16
4.4 Impacts on fauna habitats	16
4.5 Threatened species likelihood of occurrence	17
4.6 Conclusion of Assessment of Significance	17
4.7 Conclusion of EPBC assessment	17
5. Recommendations	18
6. Conclusion	19
7. Bibliography	20

Appendix A: Likelihood of occurrence	22
Appendix B: Test of significance.....	38
Appendix C: EPBC Act Significant Impact Criteria	43

List of Figures

Figure 1: The proposal overview.	3
Figure 2: The subject site and study area.....	4
Figure 3: The Biodiversity Values mapping relative to the subject site.	5
Figure 4: Vegetation communities and fauna habitats within the study area.....	14

List of Tables

Table 1: Threatened species with the potential to be affected by the proposal.....	17
--	----

Abbreviations

Abbreviation	Description
BAM	Biodiversity Assessment Method
BC Act	<i>NSW Biodiversity Conservation Act 2016</i>
BDAR	Biodiversity Development Assessment Report
CEEC	Critically Endangered Ecological Community
DCCEEW	Commonwealth Department of Climate Change, Energy, the Environment and Water
DCCEEW (NSW)	NSW Department of Climate Change, Energy, the Environment and Water
EEC	Endangered Ecological Community
ELA	Eco Logical Australia Pty Ltd
EP&A Act	<i>NSW Environmental Planning and Assessment Act 1979</i>
EPBC Act	<i>Commonwealth Environment Protection and Biodiversity Conservation Act 1999</i>
FM Act	<i>NSW Fisheries Management Act 1994</i>
FFA	Flora and Fauna Assessment
GIS	Geographic Information System
GPS	Global Positioning System
IBRA	Interim Biogeographic Regionalisation for Australia
LGA	Local Government Area
NSW	New South Wales
OEH	NSW Office of Environment and Heritage
PCT	Plant Community Type
SEPP	State Environmental Planning Policy
SSD	State Significant Development
SSI	State Significant Infrastructure
TEC	Threatened Ecological Community
VIS	Vegetation Information System
WM Act	<i>NSW Water Management Act 2000</i>

Executive Summary

This report describes the biological environment and assesses the potential effects on threatened and migratory species, endangered populations and ecological communities of a proposal to realign the Wombat Walk Connector mountain bike trail at Thredbo Alpine Resort.

The construction of the proposed realignment will require the clearing of small trees, shrubs and groundcovers in a 2.5-3 m wide corridor where the trail traverses remnant native vegetation. The trail design and construction incorporates a range of measures to minimize and mitigate the impacts on vegetation communities and fauna habitats, and on the environment generally. The alignment has been chosen to avoid locations that are particularly sensitive i.e. Subalpine Riparian Scrub and will largely be restricted to already heavily disturbed ski slopes.

The proposal will not trigger the Biodiversity Offsets Scheme (BOS), as it will not encroach on any area of land identified on the Biodiversity Values map, and the total clearing of native vegetation is significantly less than the 1 ha threshold at which the BOS is triggered.

The study area and immediate surrounds was found to support one native vegetation community; Subalpine Woodland, including shrubland derived from it, with much of the study area comprising Exotic Grassland or other heavily disturbed vegetation. No threatened flora species were recorded within the study area during the survey period and none are considered likely to occur there given the general absence of suitable habitats. The study area does not support any endangered ecological communities. Only approximately 0.01 ha of Subalpine Woodland is expected to be affected in association with the proposal. Whilst this comprises an adverse impact, it is considered acceptable given the very small proportion of the extant extent of the community within the Thredbo Resort Area, and within the locality, that will be affected.

Whilst the study area provides a small amount of potential habitat for threatened fauna species such as the Broad-toothed Rat, Eastern Pygmy-possum, Gang-gang Cockatoo, Olive Whistler, Scarlet Robin, and Flame Robin, similar habitats are extensive in the locality and the habitats to be affected are small in the context of the extent of similar habitats contiguous with the study area. Furthermore, the proposal will not affect any potentially important habitats for threatened fauna species. The proposal will not sever any linkages between habitats or otherwise permanently restrict fauna movement.

An assessment of the effects of the proposal on threatened species, populations and ecological communities which may be directly or indirectly affected by the proposal was undertaken by applying the five factors from Section 7.3 of the *Biodiversity Conservation Act 2016*. This assessment concluded that the proposal is unlikely to have a significant effect on threatened species, populations or ecological communities or their habitats.

Following consideration of the administrative guidelines for determining significance under the *Commonwealth Environment Protection & Biodiversity Conservation Act 1999*, it is concluded that the proposal is unlikely to have a significant impact on matters of National Environmental Significance or Commonwealth land, and a referral to the Commonwealth Environment Minister is not necessary.

Notwithstanding the relatively minor impacts associated with the proposal, a number of impact mitigation and amelioration measures have been recommended to be incorporated into the proposal, as identified in Section 5.

1. Introduction

Eco Logical Australia Pty Ltd (ELA) was engaged by Kosciuszko Thredbo Pty Ltd to prepare a flora and fauna assessment to accompany a proposal to modify the alignment of the Wombat Walk Connector mountain bike trail, at Thredbo Alpine Resort. This flora and fauna assessment provides the findings of a review of relevant literature, database searches and field survey. It also addresses relevant statutory considerations and makes recommendations to ameliorate the potential impacts of the proposal on vegetation and habitats.

The aim of this investigation was to assess the ecological impacts of the proposal on flora, fauna and habitats within the study area. The objectives of this investigation were:

- To identify and describe the flora species and vegetation communities present in the study area, their condition and conservation significance.
- To identify and describe the fauna habitats present in the study area and their condition.
- To identify the fauna species which are present or likely to occur in the study area, and describe their conservation significance.
- To assess the impacts of the proposal on vegetation, fauna, habitats, and other environmental features as necessary.
- To make recommendations regarding any environmental management and impact mitigation/amelioration measures, which can be implemented to limit the effects of the proposal on vegetation, fauna, habitats, and other environmental features as necessary.

1.1 The proposal

The development is for the realignment of the existing Wombat Walk Connector trail which was approved under DA 24/964. Following completion of the Alpine Coaster installation it was determined the safety clearances for the original trail sections that traverse the coaster infrastructure are not suitable. The purpose of the modification is to realign the trail away from the coaster footprint.

The proposed trail will result in an expected average disturbance footprint width of 2.5-3 m. The proposed works are expected to affect 0.01 ha of native vegetation.

The impacts of the proposed trail can be summarized as follows:

- The clearing of shrubs and groundcovers in a 2-3 m wide corridor where the trail traverses native vegetation. The disturbance corridor is required to contain the upper and lower batters and the trail surface when the trail is traversing across moderate to steep slopes. On gentler slopes the disturbance corridor will be closer to 1.5 m. The average disturbance width is expected to be 2.5 m. The clearing will be undertaken with a mix of hand tools i.e. chainsaws and brush-cutters, and machinery i.e. mini-excavator.
- In general, it is possible to align the trail to avoid tree removal, however there will be some areas where the removal of some saplings is likely to be unavoidable.
- Earthworks (cut and fill) to create the trail form. This will be undertaken with a mini-excavator.
- Importation of some decomposed granite for the track surface.

An overview of the proposal is shown in Figures 1 and further identified in Photos 1-6. A more detailed description of the proposal is also provided in the Statement of Environmental Effects for the proposal (Kosciuszko Thredbo 2024).

1.2 Direct and indirect impacts

Direct impacts on flora and fauna arising from the proposal will predominantly comprise the removal or further disturbance to approximately 100 m² of native vegetation (Subalpine Woodland and shrubland derived from the historic clearing of Subalpine Woodland).

Indirect impacts associated with the proposal are expected to be minor as:

- The footprint of the proposed direct impacts is relatively small.
- A significant proportion of the areas affected are already heavily disturbed or are on the margins of disturbed areas.
- The proposal will be implemented using low impact methods and with appropriate safeguards.

The proposal is not anticipated to result in any substantial changes in surface or subsurface hydrology which may lead to the loss or adverse modification of vegetation communities or associated habitats. Similar impacts throughout the resort and elsewhere within Kosciuszko National Park have had negligible impact on surface and subsurface hydrology, aquatic ecosystems or vegetation communities beyond the immediate footprint.

Whilst the proposal will result in increased noise and human activity whilst in operation, these increases will occur in areas that are already subject these impacts in association with chairlifts, skiing, mountain biking and resort operations. The proposal is not expected to have any substantial adverse impacts on habitat connectivity as the disruptions to existing connectivity will be minor, and only up to 3 m in width.

1.3 Subject site, study area and locality

The “subject site” comprises those areas, as described in Section 1.1 and Figures 1, which will be directly impacted by the proposal. The “study area” extends approximately 10 m beyond the limits of the subject site given the indirect impacts anticipated beyond the development footprint, as shown in Figure 2.

The locality for the purposes of this report is the area of land within a 5 km radius of the study area.




Figure 1: The proposal overview.




Figure 2: The subject site and study area.



Figure 3: The Biodiversity Values mapping relative to the subject site.



Photo 1: The proposed realignment exits the existing Wombat Walk and then stays to the north of the Alpine Coaster.



Photo 2: The proposed raalignment then winds down the Sundance ski run.



Photo 3: The proposed realignment eventually reconnects with the Village Loop trail.



Photo 4: The proposed realignment exits the existing Home Run trail and traverses the lower parts of the Lovers Leap trail.



Photo 5: The proposed realignment then heads north traversing existing shrub plantings and the bottom of a tree island before heading back to Valler Terminal along Sundance ski run.



Photo 6: The proposed alignment traverses the exotic grasslands of the lower parts of the Sundance ski run finishing at Valley Terminal.

1.4 Biodiversity Offset Scheme

The proposal does not trigger the NSW Biodiversity Offset Scheme (BOS) as:

- The proposal will not affect any land mapped within the Biodiversity Values Map as defined in the NSW Biodiversity Conservation Regulation 2017 (BC Reg), as shown in Figure 1.
- The area of native vegetation affected will be approximately 0.01 ha, which is well below the threshold (1 ha) trigger for the BOS.
- The proposal will not result in a significant impact on any threatened species, populations or communities listed under the NSW *Biodiversity Conservation Act 2016* (BC Act).

1.5 Topography, geology and soils

The study area occupies moderately sloping east facing slopes at an altitude of between approximately 1450 m and 1365 m Australian Height Datum (AHD). The study area is underlain by Silurian granodiorite (Ecology Australia 2002). Soils are likely to comprise a mix of alpine humus soils, comprising sandy clay loams. The proposed trail is within the catchment of several minor tributaries of the Thredbo River but will not traverse any mapped watercourses.

1.6 Disturbances

Much of the study area have already been disturbed in association with the existing ski runs. These disturbed areas are generally dominated by introduced grasses such as *Festuca rubra* (Red Fescue) and *Agrostis capillaris* (Browntop Bent), and a range of exotic herbs including *Acetosella vulgaris* (Sheep Sorrel), *Trifolium repens* (White Clover), *Taraxacum officinale* (Dandelion), and *Hypochaeris radicata* (Flatweed). The areas of remnant forest to be affected by the proposal are generally relatively undisturbed with only minor occurrences of cosmopolitan exotic grasses and herbs and scattered occurrences of other weeds, particularly *Rubus* spp. (Blackberry).

1.7 Planning and legislation

It is not the intention of this assessment to document all the legislation and planning instruments that are relevant to the proposal. A detailed analysis of the statutory environment is provided in the Statement of Environmental Effects for the proposal (Kosciuszko Thredbo 2024). However, the legislation and planning instruments which are relevant to the assessment of potential impacts on terrestrial flora and fauna are discussed in brief below.

1.7.1 Environmental Planning and Assessment Act 1979

The *Environmental Planning and Assessment Act 1979* (EPA Act) is the principal planning legislation for NSW, providing a framework for the overall environmental planning and assessment of development proposals. This proposal is to be assessed under Part 4 of the EPA Act. The EPA Act places a duty on the determining authority to adequately address a range of environmental matters including the maintenance of biodiversity and the likely impact to threatened species, populations and communities.

1.7.2 Biodiversity Conservation Act 2016

The *Biodiversity Conservation Act 2016* (BC Act) introduces a mandatory framework for addressing impacts on biodiversity from development and clearing, including the Biodiversity Offsets Scheme (BOS) and Biodiversity Assessment Method (BAM). The proposal will not trigger the BOS, as it will not affect any land identified on the Biodiversity Values map and the total clearing of native vegetation associated with the proposal will not exceed the 1 ha threshold which applies to the Thredbo Resort Area. As such,

a Biodiversity Development Assessment Report (BDAR) is not required and a flora and fauna assessment has been prepared. The impacts of the proposed development will be subject to a test of significance with respect to the Section 7.3 of the BC Act.

1.7.3 State Environmental Planning Policy (Precincts—Regional) 2021

State Environmental Planning Policy (Precincts—Regional) 2021 (Precincts-Regional SEPP) facilitates a planning framework for Special Activation Precincts (Precinct/s) in regional NSW, streamlining planning processes and guiding the delivery of the precincts. Chapter 4 Kosciuszko National Park and Alpine Resorts (SEPP Precincts-Regional 2021) identifies the Minister for Planning as the determining authority for development within the NSW Alpine Resorts. Precincts-Regional SEPP requires the Minister for Planning to refer for comment any development application in the Alpine Resorts to the Director General of the NSW Department of Environment and Heritage.

1.7.4 Environment Protection and Biodiversity Conservation Act 1999

The *Commonwealth Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) provides a national scheme for protecting the environment and conserving biodiversity values. Approval from the Commonwealth Minister is required under the EPBC Act if the action will, or is likely to, have a significant impact on matters considered to be of national environmental significance (MNES). MNES relevant to the proposal include species and ecological communities that are listed under the Act. The EPBC Act does not define significant impact but identifies matters that are necessary to take into consideration.

2. Methods

2.1 Database and literature review

Data gathered during all field studies and the literature review was analysed and interpreted in accordance with the provisions of legislation and planning controls pertaining to flora and fauna. Threatened and migratory species, threatened populations and threatened ecological communities (TECs) that have been recorded, or have the potential to occur within the locality have been assessed for their likelihood to inhabit the study area (Appendix A).

2.2 Field surveys

ELA conducted flora and fauna surveys within the study area and surrounds on 11 November 2024.

2.2.1 Flora surveys

A botanical survey was conducted in the study area by ELA Principal Ecologist Ryan Smithers on 11 November 2024.

2.2.1.1 *Community identification and floristic audit*

The study area was surveyed to document the flora species present, including those of conservation significance, and the location and extent of vegetation communities including any TECs encountered. A description of the vegetation was then prepared with general observations made of the wider area. The vegetation was assessed according to the floristic and structural classifications of Ecology Australia (2002) and classified to Plant Community Types (PCT).

2.2.1.2 *Targeted searches*

Specific searches for plant species of conservation significance known from the locality were conducted targeting areas of potential habitat.

2.2.1.3 *Limitations*

The floristic audit undertaken recorded as many species as possible and provides a comprehensive but not definitive species list. More species would probably be recorded during a longer survey over more seasons and years. Nevertheless, the techniques used in this investigation are considered adequate to gather the data necessary to identify potential ecological constraints to the proposal.

2.2.1.4 *Flora survey effort*

The flora survey effort employed a total of one person-hour.

2.2.2 Fauna surveys

Field investigations for fauna were conducted in conjunction with the flora surveys on 11 November 2024.

2.2.2.1 *Habitat analysis*

A description of the fauna habitats in the study area was prepared because the type of habitat in an area influences which animals occur there, as well as diversity and abundance. This habitat assessment also has an important role in predicting threatened fauna likely to occur in an area. The information collected usually includes the type of vegetation present, the presence/absence of rock habitats, tree hollows, ponds, streams, wetlands, foraging substrates and other features likely to attract threatened fauna. The

study area and immediate surrounds were traversed to identify habitat components, which were recorded and described.

2.2.2.2 Diurnal surveys

Specific searches were conducted for habitats or resources of relevance for those threatened fauna species known from subalpine and montane areas, and which might be anticipated to occur given the vegetation communities and habitats present. In particular, searches were undertaken for evidence of *Mastacomys fuscus* (Broad-toothed Rat) and for hollow-bearing trees and wombat burrows.

Opportunistic fauna surveys involved observations of animal activity, habitat surveys and searches for indirect evidence of fauna. Diurnal mammal searches were conducted in areas of potential habitat across the study area, with emphasis on searches for scats, tracks, burrows, diggings and scratchings.

2.2.2.3 Limitations

The results of fauna surveys can be optimised by conducting investigations over a long period to compensate for the effect of unfavourable weather, seasonal changes and climatic variation. In general, the longer the survey the more species will be detected. Results can also be improved by using a wide range of techniques, since some species are more likely to be detected by a particular method.

However, surveys are subject to constraints that determine the amount of time allocated, the methods used and the timing of the work. Thus, the results should be viewed in the light of these limitations. The fauna detected during the survey period are a guide to the native fauna present, but are by no means a definitive list of the species occurring in the study area. Nevertheless, the techniques used in this investigation are considered adequate to gather the data necessary to identify potential ecological constraints to the proposal.

2.2.2.4 Survey effort

The fauna survey effort employed a total of one person-hour.

3. Results

3.1 Database and literature review

Appendix A provides a list of threatened and migratory species and TECs that have been recorded from database searches within a 5 km radius of the study area. The potential for each of these species to occur in the study area and the importance of the habitats within the study area are also discussed in Appendix A, and a decision made regarding the need for further assessment in this report. Some species which are not known from montane habitats have been excluded from Appendix A.

3.2 Flora

The vegetation within the study area has been typed with reference to the classifications of Ecology Australia (2002) and into PCTs using the revised PCTs in eastern NSW, which are part of the State Vegetation Type Map (SVTM). The study area supports one native vegetation community; Subalpine Woodland, with Exotic Grassland and Disturbed Vegetation on the ski slopes, as shown in Figure 4.

3.2.1 Subalpine Woodland

Subalpine Woodland dominates the study area and surrounds, as shown in Figure 4. It is the most dominant community within the Thredbo Resort area covering an estimated 443 ha (Ecology Australia 2002). The community does not correlate well with any PCT, however is closest to PCT 3381 - *Kosciuszko Alpine Sally Woodland* and 3307 - *Kosciuszko-Namadgi Alpine Ash Moist Grassy Forest*.

The canopy is dominated by dense regrowth *Eucalyptus pauciflora* (Snow Gum) to a height of approximately 14-20 m and percent foliage cover (PFC) of up to 60%. There is a very sparse sub-canopy in places of *Acacia obliquinervia* (Mountain Hickory) to a height of 4 m and PFC of less than 1%. The understorey is dominated by *Bossiaea foliosa* (Leafy Bossiaea) to a height of up to 2.5 m with PFC of 70-90%. Other understorey shrubs which occur less frequently include *Olearia phlogopappa* (Dusty Daisy-bush), *Tasmannia xerophila* subsp. *xerophila* (Alpine Pepperbush), *Ozothamnus secundiflorus* (Cascade Everlasting), *Podocarpus lawrencei* (Mountain Plum Pine), and *Olearia megalophylla* (Large-leaf Daisy Bush).

The groundcover is typically sparse given the density of the understorey and includes patches of species such as *Poa ensiformis* (Purple-sheathed Tussock-grass), *Poa fawcettiae* (Smooth Blue Snowgrass), *Asperula gunnii* (Mountain Woodruff), *Senecio gunnii*, *Stellaria pungens*, *Dianella tasmanica* (Tasman Flax-lily), *Geranium potentilloides* var. *potentilloides*, *Acaena novae-zelandiae* (Bidgee Widgee), *Goodenia hederacea* subsp. *alpestris*, *Oxalis exilis*, *Hydrocotyle algida* and *Polystichum proliferum* (Mother Shield Fern). *Poa helmsii* (Broad-leaved Snowgrass) occurs where drainage is impeded.

Much of the subject site comprises a shrubland and or grassland that is derived from the clearing of Subalpine Woodland, as shown in Figure 4. The proposal will result in the further disturbance of approximately 0.01 ha of this derived shrubland / grassland.

3.2.2 Exotic Grassland and Disturbed Ski Slopes

The most heavily disturbed parts of the study area, i.e. the ski slopes, support exotic grassland, with occasional patches of native shrubs, grasses, and forbs. These areas are characterised by an abundance of exotic grasses and herbs, the most common species comprising *Festuca rubra*, *Agrostis capillaris*, and a range of exotic herbs including *Hypochaeris radicata*, *Acetosella vulgaris*, and *Achillea millefolium*.



Figure 4: Vegetation communities and fauna habitats within the study area.

3.3 Fauna

3.3.1 Fauna habitats

The study area contains a limited range of fauna habitats given its narrow linear nature and the dominance of the very small area of native vegetation to be affected. However, the study area is surrounded by extensive areas of native vegetation and as such, a relatively diverse range of native fauna are likely to occur there from time to time.

The Subalpine Woodland within the study area provides habitat for native birds, terrestrial and arboreal mammals, microchiropteran bats and invertebrates. In particular, it provides foraging value for birds such as *Callocephalon fimbriatum* (Gang-gang Cockatoo) and sheltering and nesting habitat for species such as *Acanthiza pusilla* (Brown Thornbill). Other threatened birds which may forage within the study area from time to time include *Petroica phoenicea* (Flame Robin), *Petroica boodang* (Scarlet Robin), and *Pachycephala olivacea* (Olive Whistler), all of which are known from the Thredbo Resort area, and in the case of the Flame Robin and Olive Whistler, are common within the Resort Area during the summer.

The Subalpine Woodlands in the Thredbo Valley are regrowth from catastrophic wildfires in 1925/26 (Banks 1986 in MGP 1996). The subsequent paucity of older or hollow-bearing trees limits sheltering habitats for hollow-dependent fauna species. No hollow-bearing trees or other nesting, breeding or roosting habitats for hollow-dependent fauna species were observed within the study area. However, some very small hollows and cavities may be present in some of the larger trees, and these may be used as shelter sites by skinks, and as nesting sites by small birds.

The study area provides a small amount of potential foraging and sheltering habitat for the Broad-toothed Rat, which is likely to be relatively widespread in the Thredbo Resort area (TAV 1997 and Green 2002). Other small mammal species such as *Cercartetus nanus* (Eastern Pygmy-possum), *Antechinus swainsonii* (Dusky Antechinus) and *Rattus fuscipes* (Southern Bush Rat) may also occur within the study area. The study area provides habitat for *Vombatus ursinus* (Common Wombat).

The rock habitats, logs and dead trees within the study area provide a limited basking and foraging resource for reptiles. The limited water habitats within the study area provide a small foraging and breeding resource for frogs such as *Crinia signifera* (Common Eastern Froglet) and possibly other species, although no frogs were calling during the survey period. The Exotic Grasslands are likely to be utilised by exotic species such as *Lepus timidus* (Brown Hare), *Oryctolagus cuniculus* (Rabbits) and *Cervus unicolour* (Sambar Deer).

4. Impact assessment

4.1 Impacts on vegetation communities

4.1.1 Subalpine Woodland

The proposal will result in the removal of approximately 100 m² of the Subalpine Woodland vegetation community, including derived shrubland, in association with the clearing of the 2.5 m wide alignment where the trail traverses intact patches of the community.

Ecology Australia (2002) estimate that there is approximately 443 ha of Subalpine Woodland within the Thredbo Resort area, and a further 184 ha within the Perisher Resort area, 183 ha at Mount Selwyn, and 5.7 ha at Charlotte Pass. The SVTM PCT map estimates that 69,187 ha or approximately 99% of the original extent of the *Kosciuszko Alpine Sally Woodland* is still extant. Furthermore, the vast majority of the occurrence of the community in NSW is within conservation reserves and in particular with Kosciuszko National Park.

In this context the loss of approximately 100 m² of Subalpine Woodland (0.002% of the extent of the community with the Thredbo Resort area) is a relatively minor and acceptable impact.

4.2 Impacts on threatened ecological communities

The study area does not support any threatened ecological communities.

4.3 Impacts on flora species of conservation significance

No threatened flora species, or flora species identified on the schedules of the Kosciuszko National Park Plan of Management (KNPPOM) (DEC 2006), were recorded within the study area during the survey period and none are expected to occur there.

4.4 Impacts on fauna habitats

Whilst the study area provides a small amount of known or potential habitat for a range of native fauna species, including threatened species, such as Broad-toothed Rat, Gang-gang Cockatoo, Olive Whistler, Scarlet Robin, and Flame Robin, similar habitats are widespread in adjacent areas, and elsewhere within the locality, and will continue to be available to these species. The impacts associated with the proposal are limited to the removal or modification of a relatively small amount of native vegetation (approximately 0.01 ha), and a few rocks, none of which provide important fauna habitats. Some sheltering and foraging habitat will be affected. However, this is a very small proportion of the sheltering and foraging habitat available in the areas immediately surrounding the study area, and the loss or modification of this habitat is not likely to adversely impact on fauna generally, or any threatened species.

The proposal will not affect any known Broad-toothed Rat nests or other important habitats for the species. No concentrations of scats or other evidence of nesting activity was detected during the survey period. Evidence of Broad-toothed Rat is widespread in the locality, and it is unlikely that a development such as proposed, would impact adversely of any individual or local population of the species. Impacts on the Eastern Pygmy-possum, if it does occur within the study area, would be minor given the narrow linear nature of the proposed vegetation clearing and the extensive surrounding forests and woodlands.

The proposal will not adversely affect the Gang-gang Cockatoo, Olive Whistler, Scarlet Robin or Flame Robin given the highly mobile nature of these species and the very small area of habitat affected relatively to the extent of similar habitat in the locality.

Under these circumstances, the impacts of the proposal on fauna habitats are relatively minor and acceptable.

4.5 Threatened species likelihood of occurrence

As a result of database searches and field surveys, the threatened species and communities identified in Table 1 are known or considered to have the potential to occur within the study area or immediate surrounds (Appendix A). The potential impact of the proposal on these entities has been assessed (Appendix C) pursuant to relevant statutory assessments.

Table 1: Threatened species with the potential to be affected by the proposal

Scientific Name	Common Name	FM Act	BC Act	EPBC Act	Occurrence
Fauna					
<i>Mastacomys fuscus</i>	Broad-toothed Rat	—	E	E	Potential
<i>Cercartetus nanus</i>	Eastern Pygmy-possum	—	V	—	Potential
<i>Callocephalon fimbriatum</i>	Gang-gang Cockatoo	—	E	E	Potential
<i>Petroica boodang</i>	Scarlet Robin	—	V	—	Potential
<i>Petroica phoenicea</i>	Flame Robin	—	V	—	Potential
<i>Pachycephala olivacea</i>	Olive Whistler	—	V	—	Potential

V = Vulnerable, E = Endangered

4.6 Conclusion of Assessment of Significance

A test of significance under Section 7.3 of the BC Act was undertaken for those threatened species known within the study area and immediate surrounds or with potential to occur there (Table 1). The outcome of the assessment was that it is highly unlikely that the proposal would significantly impact on those threatened entities assessed (Appendix B).

Recommendations have been provided in Section 5 to further ameliorate the potential impacts of the proposal.

4.7 Conclusion of EPBC assessment

An impact assessment under the EPBC Act was undertaken on threatened species known within the study area and immediate surrounds or with potential to occur there (Table 1).

The outcome of this assessment was that it is highly unlikely that the development would significantly impact on the threatened entities assessed (Appendix C). A referral to the Commonwealth under the EPBC Act is not recommended.

5. Recommendations

To further mitigate the potential impacts of the proposal, the following recommendations for impact mitigation and amelioration are suggested.

Vegetation and habitat management

1. All disturbance should be kept to the minimum required to achieve the proposal.
2. All machinery to be used during the construction phase should be limited to the existing disturbed areas and access tracks and the proposed trail alignments as far as is possible.
3. Appropriate safeguards should be in place during the proposed works to limit the potential for invasive plants or pathogens, chemicals or any other pollutants to enter the environment in association with the proposed development.
4. If any trees with a diameter at breast height (dbh) of greater than 200 mm are to be felled, then the tree must be inspected prior to felling by the KT Environmental Coordinator for any potential hollows or fissures that could potentially be used for sheltering by the Eastern Pygmy-possum.
5. If any wombat burrows need to be impacted by the proposal a wombat management plan should be developed for the proposal in consultation with NPWS.
6. Any Blackberry or other problematic weeds detected within the study area should be controlled.

Sediment control

7. Appropriate sediment control measures should be implemented prior to any construction work for the proposal and retained in place until exposed areas of soil or vegetation are stabilised and/or revegetated.
8. Drainage management and sediment control measures are to have particular regard to the prevention of any sedimentation of watercourses or vegetation communities adjoining the study area.

Rehabilitation

9. Rehabilitation activities should be consistent with the resort areas rehabilitation guidelines (NGH Environmental 2007).
10. Only weed-free straw or natural thatch/litter should be used in sediment control activities.

6. Conclusion

This report describes the biological environment and assesses the potential effects on threatened and migratory species, endangered populations and ecological communities of a proposal to realign the Wombat Walk Connector mountain bike trail at Thredbo Alpine Resort.

The study area and immediate surrounds was found to support one native vegetation community; Subalpine Woodland, with much of the study area comprising Exotic Grassland or other heavily disturbed vegetation. No threatened flora species were recorded within the study area during the survey period and none are considered likely to occur there given the general absence of suitable habitats. The study area does not support any endangered ecological communities. Only approximately 0.01 ha of Subalpine Woodland, and shrubland derived from it, is expected to be removed in association with the proposal.

Whilst the study area provides a small amount of potential habitat for threatened fauna species such as the Broad-toothed Rat, Eastern Pygmy-possum, Gang-gang Cockatoo, Olive Whistler, Scarlet Robin and Flame Robin, similar habitats are extensive in the locality and the habitats to be affected are small in the context of the extent of similar habitats contiguous with the study area. Furthermore, the proposal will not affect any potentially important habitats for threatened fauna species. The proposal will not sever any linkages between habitats or otherwise permanently restrict fauna movement.

An assessment of the effects of the proposal on threatened species, populations and ecological communities which may be directly or indirectly affected by the proposal was undertaken by applying the five factors from Section 7.3 of the *Biodiversity Conservation Act 2016*. This assessment concluded that the proposal is unlikely to have a significant effect on threatened species, populations or ecological communities or their habitats.

Following consideration of the administrative guidelines for determining significance under the Commonwealth *Environment Protection & Biodiversity Conservation Act 1999*, it is concluded that the proposal is unlikely to have a significant impact on matters of National Environmental Significance or Commonwealth land, and a referral to the Commonwealth Environment Minister is not necessary.

Notwithstanding the relatively minor impacts on vegetation and fauna habitats associated with the proposal, the impact mitigation measures described in Section 5 are also recommended to be incorporated into the proposal.

7. Bibliography

- Blakers, M., Davies, S.J.J.F., & Reilly, P.N. 1984. *The Atlas of Australian Birds*, Melbourne University Press, Melbourne.
- Briggs, J.H. & Leigh, J.D. 1996. *Rare or Threatened Australian Plants*, Australian NPWS, Canberra.
- Churchill, S. 1998. *Australian Bats*, Reed New Holland, Sydney.
- Cogger, H.G. 1996. *Reptiles and Amphibians of Australia*, Reed Books, Sydney
- Costermans, L. 1994. *Native Trees and Shrubs of South-Eastern Australia*, Lansdowne Publishing, Sydney.
- Costins, C., Gray, M., Totterdell, C., and Wimbush, D. 2000. *Kosciuszko Alpine Flora*. CSIRO Publishing, Victoria.
- Cropper, S.C. 1993, *Management of Endangered Plants*, CSIRO Publishing, Melbourne.
- Department of Environment and Conservation (DEC). 2006. *Plan of Management Kosciuszko National Park*. Department of Environment and Conservation, South Sydney.
- Ecology Australia. 2002. *Kosciuszko Resorts Vegetation Assessment*. A report for Planning NSW.
- Fisheries Scientific Committee. 2010. Proposed Determination *Aquatic Ecological Community in the Natural Drainage System of the Catchment of the Snowy River in NSW*.
- Gellie, N.J.H. 2006. Native vegetation of the southern forests: South-east Highlands, Australian Alps, South-west Slopes and South-east Corner bioregions. *Cunninghamia* 9, 219-254.
- Green, K. 2002. Selective predation on the broad-toothed rat, *Mastacomys fuscus* (Rodentia: Muridae), by the introduced red fox, *Vulpes vulpes* (Carnivora: Canidae), in the Snowy Mountains, Australia. *Austral Ecology* 27, 353–359.
- Kosciuszko Thredbo. 2024. *Statement of Environmental Effects: Wombat Walk Connector Realignment, Thredbo Alpine Resort*.
- NGH Environmental. 2007. *Rehabilitation Guidelines for the Resort Areas of Kosciuszko National Park*. A report for Parks and Wildlife Division. Department of Environment and Climate Change NSW.
- Margules Groome Poyry (MGP) PTY LTD. 1996. *Review of Environmental Factors: Easy Does It Ski Run Improvement Works*.
- Menkorst, P. Heinze, D. Broome, L. and Hynes, E. 2010. *Draft National Recovery Plan for the Mountain Pygmy-possum Burramys parvus*.
- McDougall, K.L. & Walsh, N.G. 2007. Treeless vegetation of the Australian Alps, *Cunninghamia*, vol. 10, pp. 1-57.

NSW National Parks and Wildlife Service 2001a *Approved Recovery Plan for the Threatened Alpine Flora Anemone Buttercup (Ranunculus anemoneus), Feldmark Grass (Erythranthera pumila), Raleigh Sedge (Carex raleighii) & Shining Cudweed (Euchiton nitidulus)*. NSW NPWS, Hurstville NSW.

NSW National Parks and Wildlife Service. 2001b. *Approved Recovery Plan for the Southern Corroboree Frog Pseudophryne corroboree*. NSW National Parks and Wildlife Service Hurstville.

NSW National Parks and Wildlife Service. 2002. *Approved Recovery Plan for the Mountain Pygmy Possum Burramys parvus*. NSW National Parks and Wildlife Service Hurstville.

NSW Scientific Committee. 2005. *Final Determination to list Montane Peatlands and Swamps of the New England Tableland, NSW North Coast, Sydney Basin, South East Corner, South Eastern Highlands and Australia Alps area as an endangered ecological community*.

Specht, R.L. 1970. Vegetation, in Leeper G.W. (ed), *The Australian Environment*, CSIRO Australia.

Strahan, R. 1995. *The Australian Museum Complete Book of Australian Mammals*, Cornstalk Publishing, Sydney.

Threatened Species Scientific Committee. 2009. *Listing Advice for the Alpine Sphagnum Bogs and Associated Fens Endangered Ecological Community*.

Thredbo Alpine Village (TAV). 1997. *Addendum to the Easy Does It Ski Run Improvement Works REF*.

Appendix A: Likelihood of occurrence

Summary of initial assessment to determine the likelihood of occurrence of threatened species, populations and ecological communities in the proposal site.

An assessment of likelihood of occurrence was made for threatened and migratory species identified from the database search. Additional flora species have been added where the study area is considered to provide potential habitat and additional fauna species that may inhabit the study area have also been included by correlating species habitat requirements with the existing environment. Five terms for the likelihood of occurrence of species are used in this report. This assessment was based on database or other records, presence or absence of suitable habitat, features of the study area, results of the field survey and professional judgement.

The terms for likelihood of occurrence are defined below:

- “yes” = the species was or has been observed on the site
- “likely” = a medium to high probability that a species uses the site
- “potential” = suitable habitat for a species occurs on the site, but there is insufficient information to categorise the species as likely to occur, or unlikely to occur
- “unlikely” = a very low to low probability that a species uses the site
- “no” = habitat on site and in the vicinity is unsuitable for the species.

Scientific name	Common name	FM Act	BC Act	EPBC Act	Habitat associations	Likelihood of occurrence
FLORA						
<i>Argyrotegium nitidulum</i> <i>syn. Euchiton nitidulus</i>	Shining Cudweed	-	V	V	A mat-forming silver-leaved perennial daisy growing in tall alpine herbfield or open heathland above or close to the treeline. The species is known in NSW only from the high alpine area in the vicinity of Mt Kosciuszko. The species was not observed within the study area despite good survey coverage. There is no suitable habitat for the species within the study area.	No
<i>Calotis glandulosa</i>	Mauve Burr Daisy	-	V	V	This species appears to be a coloniser of bare patches and occurs, often on roadsides, in the subalpine habitats of the Australian Alps. The species is also known from montane grasslands dominated by <i>Poa</i> species, Natural Temperate Grassland dominated by Kangaroo Grass, and Snow Gum Woodlands in the Monaro and Shoalhaven regions. Locally it is known from the Moonbah area. There is no suitable habitat for the species within the study area.	No
<i>Calotis pubescens</i>	Max Mueller's Burr-daisy	-	E	-	This species has been recorded from five sites in the Snowy Mountains of NSW (four of which, all in Kosciuszko National Park, are extant). It was first recorded in Victoria in the 19th Century but not seen again there until 2009 when a single large population was discovered south-east of Mt Hotham. It has also been detected recently in numerous locations in the northern parts of Kosciuszko National Park. It grows in subalpine treeless plains in herb-rich grassland (often dominated by <i>Poa hookeri</i>); not subject to periodic inundation. There is no suitable habitat for the species within the study area.	No
<i>Carex archeri</i>	Archer's Carex	-	E	-	This species is associated with alpine herbfield, sod tussock grassland or alpine heathland above 1000 m. There is no suitable habitat for the species within the study area.	No
<i>Carex raleighii</i>	Raleigh Sedge	-	E	-	This species is associated with bogs, alpine herbfield, sod tussock grassland, subalpine stream edges or moist alpine heathland. There is no suitable habitat for the species within the study area.	No
<i>Colobanthus curtisiae</i>	Curtis' Colobanth	-	-	V	Curtis' Colobanth is a small perennial herb growing to 40 mm high. It requires bare ground for recruitment from seed and responds well to some disturbance such as grazing. The species flowers from November to February and is largely self-pollinated. In New South Wales, the species is endemic to	No

Scientific name	Common name	FM Act	BC Act	EPBC Act	Habitat associations	Likelihood of occurrence
					Kosciuszko National Park where it occurs in subalpine / montane treeless zone below 1800 m altitude. There is no habitat for the species within the study area.	
<i>Glycine latrobeana</i>	Clover Glycine	-	CE	V	Clover Glycine is found across south-eastern Australia mainly in native grasslands and grassy woodlands, and less so in dry sclerophyll forests, woodlands. There is no suitable habitat for the species within the study area. In NSW this species is known from the Tantangara area in the northern parts of Kosciuszko National Park.	No
<i>Haloragis exalata</i> subsp. <i>exalata</i>	Square Raspwort	-	-	V	The Square Raspwort appears to be a post-disturbance coloniser, based on observations of large numbers of plants on disturbed roadsides, cleared power-line easements, and recently burnt or flooded areas. There is no suitable habitat for the species within the study area.	No
<i>Leucochrysum albicans</i> subsp. <i>tricolor</i>	Hoary Sunray	-	-	E	In NSW the Hoary Sunray occurs at relatively high elevations in woodland and open forest communities, in an area roughly bounded by Goulburn, Albury and Bega. The species has been recorded in the Yass Valley, Tumut, Upper Lachlan, Snowy River and Galong. The species is known from the South Eastern Highlands, Australian Alps and Sydney Basin bioregions. Herbarium records indicate that the taxa once occurred more widely in inland NSW, near Cobar, Dubbo, Lithgow, Moss Vale and Delegate. There is no suitable habitat for the species within the study area.	No
<i>Pimelea bracteata</i>	-	-	CE	CE	<i>Pimelea bracteata</i> occurs in wetlands and along waterways and stream edges in high altitude treeless subalpine valleys. It can also occur in wet heathland and closed heath. It is endemic to New South Wales where it is currently known from the Southern Tablelands. The main areas of occurrence are in the northern area of Kosciuszko National Park, Scabby Range Nature Reserve, neighbouring State Forests and freehold land.	No
<i>Prasophyllum bagoense</i>	Bago Leek-orchid	-	E	CE	The Bago Leek-orchid is endemic to NSW, and is currently known from a single population at McPhersons Plain, east of Tumbarumba in the Southern Tablelands. There is no suitable habitat within study area.	No
<i>Prasophyllum petilum</i>	Tarengo Leek Orchid	-	E	E	Tarengo Leek Orchid reaches to 35 cm tall. This species can be distinguished from the more common onion orchids (<i>Microtis</i> spp.) that grow in its habitat	No

Scientific name	Common name	FM Act	BC Act	EPBC Act	Habitat associations	Likelihood of occurrence
					by the pinkish-purple base to the leaf. The flowering time for this species varies from north to south. Populations around Muswellbrook and Ilford tend to flower in September, with the Boorowa and Hall populations flowering in October and the Queanbeyan area and Delegate populations in December. Annual abundance varies significantly depending on winter and early spring rainfall, biomass and potentially other variables including the severity of winter frosts. Natural populations are known from a total of five sites in NSW. These are near Boorowa, Queanbeyan area, Ilford, Delegate and a newly recognised population c.10 km west of Muswellbrook. It also occurs at Hall in the Australian Capital Territory. Grows in open sites within Natural Temperate Grassland at the Boorowa and Delegate sites. Also grows in grassy woodland in association with River Tussock <i>Poa labillardieri</i> , Black Gum <i>Eucalyptus aggregata</i> and tea-trees <i>Leptospermum</i> spp. near Queanbeyan and within the grassy groundlayer dominated by Kanagaroo Grass under Box-Gum Woodland at Ilford (and Hall, ACT).	
<i>Pterostylis oreophila</i>	Blue-tongued Orchid	-	CE	CE	In New South Wales, the Blue-tongued Greenhood is known from a few small populations within Kosciuszko National Park and a population of about 40 plants (possibly now extinct) in Bago State Forest and adjoining Crown Leases south of Tumut. It grows along sub-alpine watercourses under more open thickets of Mountain Tea-tree in muddy ground very close to water. It less commonly grows in peaty soils and sphagnum mounds. It flowers from November to January.	No
<i>Ranunculus anemoneus</i>	Anemone Buttercup	-	V	V	This perennial forb of the alpine and upper alpine zones tends to occur in areas where snow persists late into the warm season. The species is relatively common in the higher subalpine and alpine areas in the locality. This species was not observed within the study area despite good survey coverage.	No
<i>Rytidosperma pumilum</i>	Feldmark Grass	-	V	V	Feldmark Grass is limited to a tiny area of feldmark - about 3ha - of the Main Range of Kosciuszko National Park between Mt Northcote and Mt Lee. There is no suitable habitat for the species within the study area.	No
<i>Rytidosperma vickeryae</i>	Perisher Wallaby Grass	-	E	-	This perennial grass is associated with treeless subalpine streamside vegetation and has been recorded from Perisher, Betts, and Spencers Creeks	No

Scientific name	Common name	FM Act	BC Act	EPBC Act	Habitat associations	Likelihood of occurrence
					and tributaries, and Happy Jacks Plain. It is associated with bogs and sphagnum mounds. There is no suitable habitat for the species within the study area.	
<i>Thesium australe</i>	Austral Toadflax	-	V	V	This species is semi-parasitic on roots of a range of grass species, mainly Kangaroo Grass.	No
<i>Viola improcera</i>	Dwarf Violet	-	-	V	Dwarf Violet occurs at high altitudes above ca. 1300 m and up to at least 1800 m elevation in open shrubland and snow-gum woodland. The species is known to grow in open areas on or near summits and upper slopes on rocky soil and is readily seen along tracks. There are no records of the species in the locality and the habitat for the species is marginal at best. It is considered unlikely that it would occur within the study area.	Unlikely
<i>Xerochrysum palustre</i>	Swamp Everlasting	-	-	V	Grows in swamps and bogs which are often dominated by heaths. Also grows at the edges of bog margins on peaty soils with a cover of shrubs or grasses. Found in Kosciuszko National Park and the eastern escarpment south of Badja. Flowers appear from November to March. There is no suitable habitat for the species within the study area.	No
ENDANGERED ECOLOGICAL COMMUNITIES						
Montane Peatlands and Swamps of the New England Tableland, NSW North Coast, Sydney Basin, South East Corner, South Eastern Highlands and Australian Alps		-	EEC	-	The plant community characterizing this EEC is associated with accumulated peaty or organic-mineral sediments on poorly drained flats in the headwaters of streams. It occurs on undulating tablelands and plateaus, above 400-500 m elevation, generally in catchments with basic volcanic or fine-grained sedimentary substrates or, occasionally, granite. The vegetation communities within the study area do not comprise this community.	No
Alpine Sphagnum Bogs and Associated Fens		-	-	EEC	This EEC is typically found in alpine, subalpine and montane environments. It can usually be defined by the presence of sphagnum moss, even though it may sometimes only be a minor component. It is dominated by shrubs or species such as <i>Empodisma minus</i> and is found in permanently wet areas, such as along streams, valley edges, valley floors where soils are waterlogged. The vegetation communities within the study area do not comprise this community.	No

Scientific name	Common name	FM Act	BC Act	EPBC Act	Habitat associations	Likelihood of occurrence
Natural Temperate Grassland of the Southern Tablelands (NSW and ACT)		-	CEEC	EEC	This community is associated with valleys influenced by cold air drainage and open plains in the Southern Tablelands. The vegetation communities within the study area do not comprise this community.	No
White Box-Yellow Box-Blakely's Red Gum Grassy Woodland and Derived Native Grassland		-	EEC	EEC	Box Gum Woodland occurs where rainfall is between 400 and 1200 mm per annum, on moderate to highly fertile soils at altitudes of 170 m to 1200 m. It occurs in an arc along the western slopes and tablelands of the Great Dividing Range from Southern Queensland through NSW to central Victoria	No
Snowy River Aquatic Ecological Community		EEC	-	-	The bed, banks, floodplains and associated vegetation of the Snowy River and all its tributaries potentially comprise part of this EEC. The ephemeral watercourse within the study area does not comprise this EEC.	No
Disclaimer: Data extracted from the Atlas of NSW Wildlife and EPBC Act Protected Matters Report are only indicative and cannot be considered a comprehensive inventory.						
CE = Critically Endangered; E = Endangered; EEC = Endangered Ecological Community; V = Vulnerable						

FISH

<i>Maccullochella peelii</i>	Murray Cod	-	-	V	The Murray Cod utilises a diverse range of habitats from clear rocky streams, such as those found in the upper western slopes of NSW (including the ACT), to slow-flowing, turbid lowland rivers and billabongs. There is no suitable habitat within the study area.	No
<i>Macquaria australasica</i>	Macquarie Perch	-	-	E	The Macquarie Perch is a riverine, schooling species. It prefers clear water and deep, rocky holes with lots of cover. As well as aquatic vegetation, additional cover may comprise of large boulders, debris and overhanging banks (. Spawning occurs just above riffles (shallow running water). The Macquarie Perch was once widespread through the cooler upper reaches of the southern tributaries of the Murray-Darling river system in Victoria and New South Wales (Anonymous 1974; McDowall 1996), however its distribution did not usually extend to the sources of these rivers. There is no suitable habitat within the study area.	No

Scientific name	Common name	FM Act	BC Act	EPBC Act	Habitat associations	Likelihood of occurrence
<i>Prototroctes maraena</i>	Australian Grayling	-	E	V	Currently, the Australian Grayling occurs in streams and rivers on the eastern and southern flanks of the Great Dividing Range, from Sydney, southwards to the Otway Ranges of Victoria and in Tasmania. The species is found in fresh and brackish waters of coastal lagoons, from Shoalhaven River in NSW to Ewan Ponds in South Australia. It is absent from the inland Murray-Darling system (DPI 2006; McDowall 1980b). There is no suitable habitat within the study area.	No

MAMMALS

<i>Burramys parvus</i>	Mountain Pygmy-possum		E	E	This species lives only in the alpine and subalpine areas of the highest mountains of Victoria and NSW. It lives in rocky areas where boulders have accumulated below mountain peaks and is frequently associated with alpine heathlands dominated by Mountain Plum Pine. The nearest core habitats for the species are at Charlotte Pass. Given the absence of preferred sheltering or foraging habitat within the study area it is considered unlikely that the species would occur there.	Unlikely
<i>Cercartetus nanus</i>	Eastern Pygmy-possum		V	-	The Eastern Pygmy-possum is found in wet and dry eucalypt forest, subalpine woodland, coastal banksia woodland and wet heath. Pygmy-Possums feed mostly on the pollen and nectar from Banksias, Eucalypts and understorey plants and will also eat insects, seeds and fruit. The presence of Banksia sp. and Leptospermum sp. are an important habitat feature. Small tree hollows are favoured as day nesting sites, but nests have also been found under bark, in old bird nests and in the branch forks of tea-trees. The Eastern Pygmy-possum appears to be mainly solitary, each individual using several nests, with males having non-exclusive home-ranges of about 0.68 hectares and females about 0.35 hectares. They are mainly nocturnal. The Eastern Pygmy-possum is found in south-eastern Australia, from southern Queensland to eastern South Australia and in Tasmania. In NSW it extends from the coast inland as far as the Pilliga, Dubbo, Parkes and Wagga Wagga on the western slopes.	Potential
<i>Dasyurus maculatus</i>	Spotted-tailed Quoll		V	E	The species prefers moist forest types and is often associated with escarpments. There is no denning habitat for the species within the study area and the potential foraging habitat within the study area would form only	Unlikely

Scientific name	Common name	FM Act	BC Act	EPBC Act	Habitat associations	Likelihood of occurrence
					a small proportion of the home range of the species, which has been estimated at between 800 ha and 2000 ha.	
<i>Mastacomys fuscus</i>	Broad-toothed Rat		E	E	This species occurs in two widely separated areas in NSW, the Barrington Tops area and the wet alpine and subalpine heaths and woodlands of the Kosciuszko NP and adjacent areas. The species lives in a complex of runways through dense vegetation of wet grass, sedge or heath and under the snow in winter.	Potential
<i>Petauroides volans</i>	Southern Greater Glider		E	E	This species is associated with tall moist forests. It is considered unlikely that it would occur within the study area.	Unlikely
<i>Petrogale penicillata</i>	Brush-tailed Rock-wallaby		E	V	This species prefers rocky habitats, including loose boulder-piles, rocky outcrops, steep rocky slopes, cliffs, gorges and isolated rock stacks. The nearest known population is more than 50 km southeast of the study area.	No
<i>Phascolarctos cinereus</i>	Koala		V	E	Associated with both wet and dry Eucalypt forest and woodland that contains a canopy cover of approximately 10 to 70% with acceptable Eucalypt food trees. It is highly unlikely that the species would ever occur in the study area and would not be resident there.	No
<i>Pseudomys fumeus</i>	Smoky Mouse		E	E	Occurs in heath on ridge tops and slopes in sclerophyll forests, heathland and open forest along the coast and inland to sub-alpine regions. Occasionally occurs in ferny gullies. The species has recently been recorded at numerous locations in the northern parts of Kosciuszko National Park. It is considered unlikely that the species would occur within the study area or immediate surrounds give its rarity and the nature of the habitats there.	Unlikely
<i>Pteropus poliocephalus</i>	Grey-headed Flying-Fox		V	V	Inhabits a wide range of habitats including rainforest, mangroves, paperbark forests, wet and dry sclerophyll forests and cultivated areas. Camps are often located in gullies, typically close to water, in vegetation with a dense canopy. There are no camps in the locality and the species would not occur within the study area.	No
AMPHIBIANS						
<i>Litoria spenceri</i>	Spotted Tree Frog		CE	E	The Spotted Tree Frog is associated with a range of vegetation communities from montane forest at high altitudes to wet and dry forest at moderate to	No

Scientific name	Common name	FM Act	BC Act	EPBC Act	Habitat associations	Likelihood of occurrence
					low altitudes respectively. It occurs along sections of streams with steep banks, invariably in steeply dissected country or gorges with numerous rapids and waterfalls. It is restricted to riffle and cascade stream sections with exposed rock banks, resulting in a highly patchy distribution along most streams. Adults and juveniles most likely remain in the vicinity of the stream, rarely venturing far from the riparian zone. Tadpoles occur predominantly in slow-flowing sections of streams. There is no suitable habitat within the study area.	
<i>Litoria raniformis</i>	Southern Bell Frog		E	V	This species is usually found in or around permanent or ephemeral Black Box/Lignum/Nitre Goosefoot swamps, Lignum/Typha swamps and River Red Gum swamps or billabongs along floodplains and river valleys and in irrigated rice crops, particularly where there is no available natural habitat. There is no suitable habitat within the study area.	No
<i>Litoria verreauxii alpina</i>	Alpine Tree Frog		E	V	This species occurs in the alpine and subalpine zones of south-eastern NSW and Victoria. It is found in a wide variety of habitats including woodland, heath, grassland and herbfields. It breeds in natural and artificial wetlands including ponds, bogs, fens, streamside pools, dams and drainage channels that are still or slow flowing. The species has disappeared from much of its former range in the last 20 years and is restricted to a few breeding sites in murky ponds. It has recently been detected in many locations within the within the northern parts of Kosciuszko National Park suggesting a large population there. There is no suitable breeding habitat for the species within the study area and it is highly unlikely that it would occur there.	Unlikely
<i>Pseudophryne corroboree</i>	Southern Corroboree Frog		CE	CE	The Southern Corroboree Frog is limited to sphagnum bogs of the northern Snowy Mountains, in a strip from the Maragle Range in the northwest, through Mt Jagungal to Smiggin Holes in the south. Its range is entirely within Kosciuszko National Park. This species is all but extinct in the wild. It is no longer present at its former southern limit at Smiggin Holes.	No
REPTILES						
<i>Aprasia parapulchella</i>	Pink-tailed Worm Lizard		V	V	Inhabits sloping, open woodland areas with predominantly native grassy ground layers, particularly those dominated by Kangaroo Grass. Sites are typically well-drained, with rocky outcrops or scattered, partially-buried	No

Scientific name	Common name	FM Act	BC Act	EPBC Act	Habitat associations	Likelihood of occurrence
					rocks. Appear to spend considerable time in burrows below rocks; the burrows have been constructed by and are often still inhabited by small black ants and termites. The study area does not support suitable habitat for the species. The nearest records of the species are more than 50 km away at Cooma.	
<i>Cyclodomorphus praealtus</i>	Alpine She-oak Skink		E	E	In NSW, the species is known from open alpine heath and tussock grassland within the Kosciuszko region, preferring treeless or lightly treed areas. The study area does not include any suitable habitat for this species and it is considered unlikely that it would occur there.	Unlikely
<i>Eulamprus kosciuskoi</i>	Alpine Water Skink		-	E	The alpine water skink has been observed at high elevations (above 1000 m above sea level (asl) and up to 2000 m asl), basking on granite boulders, tussocks and sphagnum moss along small alpine streams, bogs, and wet heath. It has also been observed along drainage lines, in tussock grasses, and up to hundreds of metres upslope, or otherwise away, from typical wetter habitat. In the Snowy Mountains and Brindabella Range, the species exhibits a high degree of habitat specialisation, being confined to sphagnum bog, fen, wet heath and, less frequently, wet sod-tussock grassland. The study area is typically too densely treed to provide suitable habitat for the species which is associated with more open and wet habitats at higher altitudes.	No
<i>Liopholis guthega</i>	Guthega Skink		E	E	This species is known from the Snowy Mountains and the Bogong High Plains and is associated with rocky areas in a range of alpine and subalpine vegetation communities. The species lives in extensive colonies associated with a deep burrow network that is constructed in eroded granite and humus soils beneath boulders and shrubs. The species is not associated with Subalpine Woodland and is highly unlikely to occur within the study area.	No
<i>Liopholis montana</i>	Mountain Skink		E	E	This species is known from the Snowy Mountains, Victorian Alps and ACT and is associated with rocky areas in a range of alpine and subalpine and montane vegetation communities above 600 m. The species lives in extensive colonies associated with a deep burrow network that is constructed in eroded granite and humus soils beneath boulders and shrubs. The study area does not include any suitable habitat for this species. The study area is typically too densely treed to provide suitable habitat for the	Unlikely

Scientific name	Common name	FM Act	BC Act	EPBC Act	Habitat associations	Likelihood of occurrence
					species which is associated with more open and typically rocky habitats. There are no records within the Thredbo resort area, with the nearest records being at Blue Cow mountain.	
<i>Pseudemoia cryodroma</i>	Alpine Bog Skink		-	E	The alpine bog skink occurs primarily in alpine bog, riparian and wet heath areas above 1100 m elevation, and less commonly in alpine and subalpine grassland and dry treeless heath, drainage lines in subalpine meadows, and in snow gum woodland. The study area is typically too densely treed to provide suitable habitat for the species which is associated with more open and wet habitats.	No
BIRDS						
<i>Anthochaera phrygia</i>	Regent Honeyeater		CE	CE, M	Associated with temperate eucalypt woodland and open forest including forest edges, wooded farmland and urban areas with mature eucalypts, and riparian forests of River Oak (<i>Casuarina cunninghamiana</i>). The Regent Honeyeater primarily feeds on nectar from box and ironbark eucalypts and occasionally from banksias and mistletoes. As such it is reliant on locally abundant nectar sources with different flowering times to provide a reliable supply of nectar. The species would not occur within the study area.	No
<i>Artamus cyanopterus cyanopterus</i>	Dusky Woodswallow		V	-	Dusky woodswallows are widespread in eastern, southern and south western Australia. The species occurs throughout most of New South Wales, but is sparsely scattered in, or largely absent from, much of the upper western region. Most breeding activity occurs on the western slopes of the Great Dividing Range. They primarily inhabit dry, open eucalypt forests and woodlands, including mallee associations, with an open or sparse understorey of eucalypt saplings, acacias and other shrubs, and ground-cover of grasses or sedges and fallen woody debris. It has also been recorded in shrublands, heathlands and very occasionally in moist forest or rainforest. Also found in farmland, usually at the edges of forest or woodland.	Unlikely
<i>Botaurus poiciloptilus</i>	Australasian Bittern		V	E	This species favours permanent freshwater wetlands with tall, dense vegetation, particularly bulrushes and spikerushes. It hides during the day amongst dense reeds and feeds at night. It breeds during summer with nest built in secluded places in densely vegetated wetlands on a platform of reeds. There is no habitat for the species within the study area.	No

Scientific name	Common name	FM Act	BC Act	EPBC Act	Habitat associations	Likelihood of occurrence
<i>Callocephalon fimbriatum</i>	Gang-gang Cockatoo		E	E	Gang-gang Cockatoos live as pairs inhabiting woodlands of south-eastern Australia. The species feeds primarily on the seeds of eucalypts and acacias and breeds in tree hollows. The species is typically associated with taller montane forests in the region but is sometimes observed foraging in Snow Gums and on the side of roads. It's likely that the species would forage within the study area from time to time.	Potential
<i>Calyptorhynchus lathamii</i>	South-eastern Glossy Black-Cockatoo		V	V	This species occurs in forests and woodlands where She-oak feeding resources are prevalent and large tree hollows exist for breeding. There is no foraging or breeding habitat for the species within the study area and it is highly unlikely that it would occur there.	Unlikely
<i>Climacteris picumnus</i>	Brown Treecreeper (eastern subspecies)		V	—	This species occurs in Eucalypt forests and woodlands of inland plains and slopes of the Great Dividing Range. It is not usually found in woodlands with a dense shrub layer. Fallen timber is an important habitat component for foraging. It is a sedentary species and is resident in many locations throughout its range, and is present in all seasons or year-round at many sites. Hollows in standing dead or live trees and tree stumps are essential for nesting. There is no habitat for the species within the study area.	No
<i>Daphoenositta chrysoptera</i>	Varied Sittella		V	—	The Varied Sittella is sedentary and inhabits most of mainland Australia except the treeless deserts and open grasslands, with a nearly continuous distribution in NSW from the coast to the far west. It inhabits eucalypt forests and woodlands, especially rough-barked species and mature smooth-barked gums with dead branches, mallee and Acacia woodland. It is considered unlikely that the species would occur within the study area.	Unlikely
<i>Falco hypoleucos</i>	Grey Falcon		E	-	The Grey Falcon is sparsely distributed in NSW, chiefly throughout the Murray-Darling Basin, with the occasional vagrant east of the Great Dividing Range. The breeding range has contracted since the 1950s with most breeding now confined to arid parts of the range. There are possibly less than 5000 individuals left. Population trends are unclear, though it is believed to be extinct in areas with more than 500mm rainfall in NSW. The Grey Falcon is sparsely distributed in NSW, chiefly throughout the Murray-Darling Basin, with the occasional vagrant east of the Great Dividing Range. The breeding range has contracted since the 1950s with most breeding now	No

Scientific name	Common name	FM Act	BC Act	EPBC Act	Habitat associations	Likelihood of occurrence
					confined to arid parts of the range. There are possibly less than 5000 individuals left. Population trends are unclear, though it is believed to be extinct in areas with more than 500mm rainfall in NSW.	
<i>Grantiella picta</i>	Painted Honeyeater		V	V	The Painted Honeyeater is a nomadic species that occurs predominantly on the inland slopes of the Great Dividing Range. It inhabits Boree (<i>Acacia pendula</i>), Brigalow (<i>A. harpophylla</i>) and Box-Gum Woodlands and Box-Ironbark Forests. It is a specialist feeder on the fruits of mistletoes growing on woodland eucalypts and acacias, preferring mistletoes of the genus <i>Amyema</i> . Nesting occurs from spring to autumn in a small, delicate nest hanging within the outer canopy of drooping <i>Eucalyptus</i> spp., <i>Allocasuarina</i> and <i>Casuarina</i> spp. (Sheoaks), <i>Melaleuca</i> sp. (Paperbark) or Mistletoe branches. It is highly unlikely that the species would occur within the study area.	Unlikely
<i>Lathamus discolor</i>	Swift Parrot		CE	CE	Breeds in Tasmania between September and January. Migrates to mainland in autumn, where it forages on profuse flowering Eucalypts. Hence, in this region, autumn and winter flowering eucalypts are important for this species. Favoured feed trees include winter flowering species such as Swamp Mahogany (<i>Eucalyptus robusta</i>), Spotted Gum (<i>Corymbia maculata</i>), Red Bloodwood (<i>C. gummifera</i>), Mugga Ironbark (<i>E. sideroxylon</i>), and White Box (<i>E. albens</i>). It is considered highly unlikely that the species would occur within the study area.	Unlikely
<i>Neophema chrysogaster</i>	Orange-bellied Parrot		E	CE, M	Breeds only in coastal south-west Tasmania and spends the winter in coastal Victoria and South Australia. It nests in hollows in eucalypt trees which grow adjacent to its feeding plains. In early October the birds arrive in the south west and depart after the breeding season usually in March and April. It feeds on the seeds of several sedges and heath plants, including buttongrass. Its main food preferences are found in sedgelands which have not been burned for between 3-15 years. Also included in the diet are seeds of three <i>Boronia</i> species and the everlasting daisy (<i>Helichrysum pumilum</i>). After breeding, migrating birds move gradually northwards up the west coast, through the Hunter Group and King Island in Bass Strait and on to the mainland. On the journey the birds usually feed on beach-front vegetation including salt tolerant species such as sea rocket (<i>Cakile maritima</i>). They also	No

Scientific name	Common name	FM Act	BC Act	EPBC Act	Habitat associations	Likelihood of occurrence
					eat various coastal native and introduced grasses. There is no habitat for the species within the study area.	
<i>Pachycephala olivacea</i>	Olive Whistler		V	-	This species is usually associated with moist tall forests at high elevations but has been occasionally recorded at lower altitudes. Breeding occurs above 300m within habitats providing both a thick understorey and moderate canopy. In the alps the species is more typically associated with subalpine woodlands with a heathy understorey. It is likely that the species would occur within the study area from time to time.	Potential
<i>Petroica rodinogaster</i>	Pink Robin		V	-	The Pink Robin is found in Tasmania and the uplands of eastern Victoria and far south-eastern NSW, almost as far north as Bombala. It inhabits rainforest and tall, open eucalypt forest, particularly in densely vegetated gullies. In the alps the species is more typically associated with subalpine woodlands with a heathy understorey and Montane Forests rather than alpine heaths. The species may forage within the study area from time to time.	Unlikely
<i>Petroica boodang</i>	Scarlet Robin		V	-	This species is found in south-eastern Australia and south-west Western Australia. In NSW it occupies open forests and woodlands from the coast to the inland slopes. The Scarlet Robin breeds in drier eucalypt forests and temperate woodlands, often on ridges and slopes, within an open understorey of shrubs and grasses. Abundant logs and coarse woody debris are important structural components of its habitat. In autumn and winter it migrates to more open habitats such as grassy open woodland or paddocks with scattered trees. The habitat for the species within the study area is marginal however the species is recorded within the Thredbo Valley from time to time and has been recorded by the author in similar habitats to those within the study area.	Potential
<i>Petroica phoenicea</i>	Flame Robin		V	-	The Flame Robin is found in south-eastern Australia (Queensland border to Tasmania, western Victoria and south-east South Australia). In NSW it breeds in upland moist eucalypt forests and woodlands, often on ridges and slopes, in areas of open understorey. It migrates in winter to more open lowland habitats such as grassland with scattered trees and open woodland on the inland slopes and plains. The species is well known from the locality	Known

Scientific name	Common name	FM Act	BC Act	EPBC Act	Habitat associations	Likelihood of occurrence
<i>Pycnoptilus floccosus</i>	Pilotbird		V	V	and would likely use the more open habitats within the study area from time to time for foraging. The Pilotbird occurs in temperate wet sclerophyll forests and occasionally temperate rainforest, where there is dense undergrowth with abundant debris. Pilotbirds are known from the Thredbo Resort area however the species is considered unlikely to occur within the study area given the open nature of the habitats there..	Unlikely
<i>Rostratula australis</i>	Australian Painted Snipe		E	E	In NSW, records of the Painted Snipe are from the Murray-Darling Basin, including the Paroo wetlands, Lake Cowal, Macquarie Marshes, Fivebough Swamp, and swamps near Balldale and Wanganella. Other important locations with recent records include wetlands on the Hawkesbury River and the Clarence and lower Hunter Valleys. It prefers the fringes of swamps, dams and nearby marshy areas, where there is a cover of grasses, Lignum, low scrub or open timber. It nests on the ground amongst tall vegetation, such as grasses, tussocks or reeds. There is no suitable habitat for the species within the study area.	No
MIGRATORY TERRESTRIAL AND WETLAND SPECIES LISTED UNDER EPBC ACT						
<i>Hirundapus caudacutus</i>	White-throated Needletail		—	V, M	Forages aerially over a variety of habitats usually over coastal and mountain areas, most likely with a preference for wooded areas. Has been observed roosting in dense foliage of canopy trees, and may seek refuge in tree hollows in inclement weather.	Unlikely
<i>Merops ornatus</i>	Rainbow Bee-eater		—	M	Resident in coastal and subcoastal northern Australia; regular breeding migrant in southern Australia, arriving September to October, departing February to March, some occasionally present April to May. Occurs in open country, chiefly at suitable breeding places in areas of sandy or loamy soil: sand-ridges, riverbanks, road-cuttings, sand-pits, occasionally coastal cliffs (ibid). Nest is a chamber at the end of a burrow, up to 1.6 m long, tunnelled in flat or sloping ground, sandy back or cutting (ibid). The species would not occur within the study area.	No

Scientific name	Common name	FM Act	BC Act	EPBC Act	Habitat associations	Likelihood of occurrence
<i>Monarcha melanopsis</i>	Black-faced Monarch		—	M	This migratory species is known to breed in damp forest types and forage in rainforest and eucalypt forest. The species would not occur within the study area.	No
<i>Myiagra cyanoleuca</i>	Satin Flycatcher		—	M	This species inhabits lowland eucalypt forests. It is known to nest in dense gully vegetation. The species would not occur within the study area.	No
<i>Neophema chrysogaster</i>	Orange-bellied Parrot		E	E, M	SEE DIURNAL BIRDS ABOVE	No
<i>Rhipidura rufifrons</i>	Rufous Fantail		—	M	This migratory species forages by catching flying insects and is known to utilise the aerial foraging space above the dense understorey in damp forests or beside rivers. The species would not occur within the study area.	No
<i>Xanthomyza phrygia</i>	Regent Honeyeater		E	E, M	SEE DIURNAL BIRDS ABOVE	No
<i>Gallinago hardwickii</i>	Latham's Snipe		E	M	Resides in swamps, dams and nearby marshy areas that contain grasses, lignum, low scrub or open timber that provides cover. It is considered highly unlikely that the species would occur within the study area.	Unlikely
<i>Motacilla flava</i>	Yellow Wagtail		—	M	Frequents open wetlands along the bare shores of freshwater swamps, crops and bare bore drains, as well as short-grassed fields and rocky coasts. It is considered highly unlikely that the species would occur within the study area.	Unlikely

Disclaimer: Data extracted from the Atlas of NSW Wildlife and EPBC Act Protected Matters Report are only indicative and cannot be considered a comprehensive inventory. 'Migratory marine species' and 'listed marine species' listed on the EPBC Act (and listed on the protected matters report) have not been included in this table, since they are considered unlikely to occur within the study area due to the absence of marine and wetland habitats.

CE = Critically Endangered; E = Endangered; V = Vulnerable; M = Migratory

Appendix B: Test of significance

Test of significance pursuant to section 7.3 of the BC Act: Five-part test

An assessment of the effects of the proposal on threatened species, populations and ecological communities which may be directly or indirectly affected by the proposal may be carried out by applying the five factors from Section 7.3 of the BC Act.

This test of significance is presented below for the threatened fauna species:

- *Mastacomys fuscus* (Broad-toothed Rat)
- *Cercartetus nanus* (Eastern Pygmy-possum)
- *Callocephalon fimbriatum* (Gang-gang Cockatoo)
- *Petroica phoenicea* (Flame Robin)
- *Petroica boodang* (Scarlet Robin)
- *Pachycephala olivacea* (Olive Whistler)

(a) in the case of a threatened species, whether the proposed development or activity is likely to have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction,

Endangered Species

Fauna

Broad-toothed Rat *Mastacomys fuscus* (potential occurrence).

The Broad-toothed Rat generally occurs in two widely separated areas in NSW, the Barrington Tops area and the wet alpine and subalpine heaths and woodlands of the Kosciuszko NP and adjacent areas. The species lives in a complex of runways through dense vegetation of wet grass, sedge or heath and under the snow in winter. Home range size is thought to range between approximately 0.1 ha and 0.27 ha. Individuals nest alone over summer but congregate in communal nests during winter. The species is thought to be locally common in the alpine and high subalpine tracts of the Snowy Mountains area (Green 2002), where suitable habitats are present.

The study area provides a small amount of marginal potential foraging and sheltering habitat for the Broad-toothed Rat. Whilst no evidence of the species was observed within the study area, it has been observed nearby, and it is possible that the species occurs within the study area from time to time.

The proposed development will affect only a very small amount of the potential habitat for the species in the Thredbo Resort area and will not affect any key resources for the species. As such, the proposed development is unlikely to adversely affect a significant proportion of the home range of one or more Broad-toothed Rat individuals.

The proposed development will not result in habitat fragmentation which could isolate individuals or a population of the Broad-toothed Rat, given the narrowness of the clearing required. Under these circumstances, the proposed development is considered unlikely to disrupt the life cycle of the Broad-toothed Rat such that a viable local population is likely to be placed at risk of extinction.

Gang-gang Cockatoo *Callocephalon fimbriatum* (potential occurrence).

In New South Wales, the Gang-gang Cockatoo is distributed from the south-east coast to the Hunter region, and inland to the central and southern tablelands and south-west slopes. In summer, this species is generally found in tall mountain forests and woodlands, particularly in heavily timbered and mature wet sclerophyll forests. In winter, the Gang-gang Cockatoo may occur at lower altitudes in drier more open eucalypt forests and woodlands, and is often found in urban areas. It may also occur in sub-alpine Snow Gum woodland and occasionally in temperate rainforests.

The species is regularly observed at Thredbo in montane and subalpine areas. Whilst the species may forage within the study area, it would not breed there given the absence of suitable nesting habitat. Given the extensive forests within the locality, breeding and roosting habitat is likely to be relatively abundant.

The study area provides a very small area of suitable foraging resources for the species. The foraging resources (generally eucalypt trees) to be removed in association with the proposed development would not be important for the species, given the extent of foraging resources in the Thredbo Resort area.

Under these circumstances, the proposed development will not disrupt the life cycle of the Gang-gang Cockatoo such that a viable local population of the species is likely to be placed at risk of extinction.

Vulnerable Species**Fauna****Eastern Pygmy-possum *Cercartetus nanus* (potential occurrence).**

The Eastern Pygmy-possum is found in wet and dry eucalypt forest, subalpine woodland, coastal banksia woodland and wet heath. Pygmy-Possums feed mostly on the pollen and nectar from Banksias, Eucalypts and understorey plants and will also eat insects, seeds and fruit. The presence of Banksia sp. and Leptospermum sp. are an important habitat feature. Small tree hollows are favoured as day nesting sites, but nests have also been found under bark, in old bird nests and in the branch forks of tea-trees. The Eastern Pygmy-possum appears to be mainly solitary, each individual using several nests, with males having non-exclusive home-ranges of about 0.68 hectares and females about 0.35 hectares. They are mainly nocturnal. The Eastern Pygmy-possum is found in south-eastern Australia, from southern Queensland to eastern South Australia and in Tasmania. In NSW it extends from the coast inland as far as the Pilliga, Dubbo, Parkes and Wagga Wagga on the western slopes.

There are a few records of the species from Kosciuszko National Park, mainly from lower altitudes, however the species has been recorded at 1,800 m. It is likely that the Eastern Pygmy-possum occurs in the subalpine and montane habitats of the Thredbo Resort.

The proposal is unlikely to adversely affect a significant proportion of the home range of any individual Eastern Pygmy-possum given that it comprises a narrow linear development. It is unlikely that any individual Eastern Pygmy-possum would be directly affected by the proposed trail given the relatively small area to be affected during the construction phase, the fact that no hollow-bearing trees will be removed and that the disturbances during construction are likely to encourage any individuals that may be within the disturbance corridor, to move away. Direct impacts during the use of the trail are unlikely given that the species is primarily nocturnal.

The proposal is highly unlikely to disrupt the life cycle of the Eastern Pygmy-possum such that a viable local population of the species is likely to be placed at risk of extinction.

Flame Robin *Petroica phoenicea* (known occurrence).

The Flame Robin is found in south-eastern Australia (Queensland border to Tasmania, western Victoria and south-east South Australia). In NSW it breeds in upland moist eucalypt forests and woodlands, often on ridges and slopes, in areas of open understorey. It migrates in winter to more open lowland habitats such as grassland with scattered trees and open woodland on the inland slopes and plains. There are numerous records of the species throughout the NSW Alps, and the species was observed in the study area during the survey period. It is well known from the Thredbo Resort area and is one of the most common birds of open habitats outside of the winter period.

The proposal will affect a very small amount of potential nesting and foraging habitat for the species. This is negligible in the context of the extensive areas of similar habitat within the Thredbo Resort area that will not be affected by the proposed development and which will continue to be available to the species. The species is not sedentary and undertakes substantial seasonal migrations, reducing the species dependence on any specific area of known or potential habitat.

Under these circumstances, the proposed development is unlikely to disrupt the life cycle of the Flame Robin such that a viable local population of the species is likely to be placed at risk of extinction.

Scarlet Robin *Petroica boodang* (potential occurrence).

This Scarlet Robin is found in south-eastern Australia and south-west Western Australia. In NSW it occupies open forests and woodlands from the coast to the inland slopes. The Scarlet Robin breeds in drier eucalypt forests and temperate woodlands, often on ridges and slopes, within an open understorey of shrubs and grasses. Abundant logs and coarse woody debris are important structural components of its habitat. In autumn and winter it migrates to more open habitats such as grassy open woodland or paddocks with scattered trees. The habitat for the species within the study area is marginal however the species is recorded within the Thredbo Valley from time to time.

The proposal will affect a very small amount of potential nesting and foraging habitat for the species. This is negligible in the context of the extensive areas of similar habitat within the Thredbo Resort area that will not be affected by the proposed development and which will continue to be available to the species. The species is not sedentary and undertakes substantial seasonal migrations, reducing the species dependence on any specific area of known or potential habitat.

Under these circumstances, the proposed development is unlikely to disrupt the life cycle of the Scarlet Robin such that a viable local population of the species is likely to be placed at risk of extinction.

Olive Whistler *Pachycephala olivacea* (potential occurrence).

The Olive Whistler is found in south-eastern Australia (Queensland border to Tasmania, western Victoria and south-east South Australia). In the NSW Alps, it is associated with areas of tall dense heath, particularly riparian Tea-tree scrubs. It breeds in the thick understorey of moist eucalypt forests and subalpine woodlands. It migrates in winter to lowland habitats. There are numerous records of the species throughout the NSW Alps including within the Thredbo Resort area where it is considered a common resident. The species was not recorded within the study area during the survey period however it is likely to occur there from time to time.

The proposed development will result in the loss of a small amount of potential foraging and breeding habitat for the Olive Whistler. Whilst this comprises an adverse impact on the species, the habitat to be removed is very small relative to the extensive areas of similar habitat which occurs within the Thredbo

Resort area and elsewhere in the locality. Extensive areas of potential habitat for the species is contiguous with the study area in the extensive Subalpine Woodland and Subalpine Riparian Scrub within the Thredbo Valley. The species is highly mobile and considered to be common within the Thredbo Valley and the Thredbo Valley population is considered to be contiguous with other populations to the north and south (MGP 1996).

Under these circumstances it is considered unlikely that the proposed development would affect the life cycle of the Olive Whistler such that a viable local population of the species is likely to be placed at risk of extinction.

(b) in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:

(i) is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or

(ii) is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction,

There are no endangered or critically endangered ecological communities within the study area.

(c) in relation to the habitat of a threatened species or ecological community:

(i) the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and

The proposed development will impact on only a very small area (0.01 ha) of potential habitat for the Broad-toothed Rat and Eastern Pygmy-possum and will not affect any known Broad-toothed Rat communal nesting or likely breeding sites for either species. The proposed development will result in the modification of a very small amount of potential foraging and breeding habitat (0.01 ha) for the Flame Robin, Scarlet Robin and Olive Whistler, and only a very small amount of potential foraging habitat for the Gang-gang Cockatoo.

(ii) whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed development or activity, and

The proposed development primarily involves a narrow band of clearing of understorey and groundcover vegetation. The proposed clearing will not sever connectivity between the fauna habitats within the study area and contiguous habitats, or isolate any fauna populations which may occur within the study area. The disruptions to connectivity between fauna habitats will be minor, typically 2.5 m in width. This is considered highly unlikely to sever connectivity between habitats even for relatively immobile species with small home ranges such as some small mammals and reptiles.

The effects of the action proposed on habitat connectivity will be minor and the native fauna which may occur within the study area from time to time, will continue to be able to traverse the study area.

(iii) the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality,

The potential Broad-toothed Rat habitats to be affected comprise a small area of marginal habitat relative to the extensive areas of similar and superior habitats provided by contiguous vegetation. The alpine, subalpine and montane heaths in the locality provide superior habitat for the species than the habitats within the study area. No evidence of any important communal nesting sites was observed within the study area. Under these circumstances, the habitats to be affected are not considered to be particularly important for Broad-toothed Rat.

The habitat to be removed by the proposal is highly unlikely to be important to the long-term survival of the Eastern Pygmy-possum in the locality given that it comprises only a relatively small amount of potential habitat for the species relative to the extensive areas of remnant forest, woodland and heath within the locality.

In the context of the extent of similar habitat available for the Gang-gang Cockatoo, Olive Whistler, Scarlet Robin and Flame Robin in the Thredbo Resort area and elsewhere in the locality, the habitats within the study area are not considered to be important.

d) whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly),

The proposed development will not affect any area of outstanding biodiversity value.

(e) whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key threatening process.

The proposed development will remove 0.01 ha of remnant native vegetation. Whilst this constitutes the Key Threatening Process 'Clearing of native vegetation', the contribution to this key threatening process is relatively minor considering the extent of remnant forest in the locality and the extant extent of the vegetation communities that will be affected.

Appendix C: EPBC Act Significant Impact Criteria

The EPBC Act Administrative Guidelines on Significance set out ‘**Significant Impact Criteria**’ that are to be used to assist in determining whether a proposed action is likely to have a significant impact on matters of national environmental significance. Matters listed under the EPBC Act as being of national environmental significance include:

- Listed threatened species and ecological communities;
- Listed migratory species;
- Wetlands of International Importance;
- The Commonwealth marine environment;
- World Heritage properties;
- National Heritage places;
- Nuclear actions; and
- Great Barrier Reef.

Specific ‘**Significant Impact Criteria**’ are provided for each matter of national environmental significance except for threatened species and ecological communities in which case separate criteria are provided for species listed as endangered and vulnerable under the EPBC Act.

Threatened and migratory species listed under the EPBC Act that are considered likely or potentially to occur within the study area are given in **Appendix A** of the Report. The only Commonwealth listed species which are considered to have the potential to occur within the study area are the Broad-toothed Rat and the Gang-gang Cockatoo.

The relevant Significant Impact Criteria have been applied to determine the significance of impacts associated with the proposal.

Matters to be addressed	Impact
(a) any environmental impact on a World Heritage Property or National Heritage Places;	No. The proposal does not impact on a World Heritage Property or a National Heritage Place as addressed in the SEE. (listed natural: Australian Alpine National Parks and Reserves; nominated historic: Snowy Mountains Scheme NSW).
(b) any environmental impact on Wetlands of International Importance;	No. The proposal will not affect any part of Ramsar wetland.
(c) any impact on Commonwealth Listed Critically Endangered or Endangered Species;	Yes. The study area provides potential habitat for two Commonwealth listed endangered species: the Gang-gang Cockatoo and the Broad-toothed Rat. The significant impact criteria for endangered species are discussed below: a. lead to a long-term decrease in the size a population of a species, The proposed action will only affect a very small amount of potential habitat for these species in the context of the extent of potential habitat in the locality. The proposal will not affect any breeding or roosting habitat or otherwise adversely impact these species.

Matters to be addressed	Impact
	<p>Under these circumstances, it is considered highly unlikely that the proposed action will lead to a long-term decrease in the size of the Gang-gang Cockatoo or Broad-toothed Rat population.</p>
	<p>b. reduce the area of occupancy of the species</p> <p>The proposed action will be limited to the loss or further modification of 0.01 ha of native vegetation which is a small amount of habitat in the context of the extent of similar habitats in the locality generally. The proposed works will not affect any key habitat resources for the Gang-gang Cockatoo or Broad-toothed Rat; nor affect these species ability to access habitats within or beyond the study area.</p> <p>Under these circumstances, the proposed action is highly unlikely to reduce the area of occupancy of the local population of the Gang-gang Cockatoo or Broad-toothed Rat.</p>
	<p>c. fragment an existing population into two or more populations</p> <p>The proposed action will be limited to the loss or further modification of 0.01 ha of native vegetation which is a small amount of habitat in the context of the extent of similar habitats in the locality generally. The proposed works will not affect any key habitat resources for the Gang-gang Cockatoo or Broad-toothed Rat; nor affect these species ability to access habitats within or beyond the study area.</p> <p>Under these circumstances, the proposed action will not fragment an existing population of the Gang-gang Cockatoo or Broad-toothed Rat into two or more populations.</p>
	<p>d. adversely affect habitat critical to the survival of a species</p> <p>No habitat within the development site is considered likely to be critical to the survival of the Gang-gang Cockatoo or Broad-toothed Rat. There are thousands of hectares of similar habitats in the alpine and subalpine zones of the Australian alps, including elsewhere within the Thredbo Resort area. The Gang-gang Cockatoo and Broad-toothed Rat continue to occur within the Thredbo Resort Area despite a long history of similar and more extensive disturbances.</p>
	<p>e. disrupt the breeding cycle of a population</p> <p>It is considered highly unlikely that the Gang-gang Cockatoo would breed within the study area given the absence of hollow-bearing trees. Similarly, it is considered unlikely that the Broad-toothed Rat would breed within the study area given the absence of the species preferred wet heath and bog habitats.</p> <p>Under these circumstances, the proposed action will not disrupt the breeding cycle of a population of the Gang-gang Cockatoo or Broad-toothed Rat.</p>
	<p>f. modify, destroy, remove or isolate or decrease the availability or quality of habitat to the extent that the species is likely to decline</p> <p>The proposed action will modify a very small area of habitat for the Gang-gang Cockatoo and Broad-toothed Rat, but this area is unlikely to be important to these species in the context of the extent of potential habitat in the locality.</p> <p>Under these circumstances it is highly unlikely that the proposed action would modify- destroy- remove or isolate or decrease the availability or quality of habitat to the extent that the Gang-gang Cockatoo or Broad-toothed Rat is likely to decline.</p>
	<p>g. result in invasive species that are harmful to an endangered species becoming established in the endangered or critically endangered species' habitat</p> <p>The proposed action is unlikely to result in invasive species that are harmful becoming established in potential habitat of the Gang-gang Cockatoo or Broad-toothed Rat. Species such</p>

Matters to be addressed	Impact
	<p>as cats or foxes are already present in the landscape and are subject to control programs within the resort.</p> <p>h. introduce disease that may cause the species to decline</p> <p>The proposed action is unlikely to introduce disease that may cause the Gang-gang Cockatoo or Broad-toothed Rat to decline.</p> <p>i. interfere substantially with the recovery of the species.</p> <p>As the proposed action is not considered to decrease or fragment any existing populations the recovery of the Gang-gang Cockatoo or Broad-toothed Rat is unlikely to be adversely impacted.</p>
(d) any impact on Commonwealth Listed Vulnerable Species;	No. The study area does not provide any potential habitat for any Commonwealth listed vulnerable species.
(e) Any impact on a Commonwealth Endangered Ecological Community	No. The proposal will not impact any Commonwealth listed endangered ecological communities.
(f) any environmental impact on Commonwealth Listed Migratory Species;	No. The proposal will not have any adverse impacts on any listed migratory species.
(g) does any part of the Proposal involve a Nuclear Action;	No. The project does not include a Nuclear Action.
(h) any environmental impact on a Commonwealth Marine Area;	No. There are no Commonwealth Marine Areas within the study area.
(i) In addition, any direct or indirect impact on Commonwealth lands	No. The project does not directly or indirectly affect Commonwealth land.



Appendix B AHIMS Search Results

Kosciusko Thredbo

Date: 20 November 2024

Attention: Jocelyn Best

Email: jocelyn_best@evt.com

Dear Sir or Madam:

AHIMS Web Service search for the following area at Lat, Long From : -36.5041, 148.2994 - Lat, Long To : -36.4998, 148.3071, conducted by Jocelyn Best on 20 November 2024.

The context area of your search is shown in the map below. Please note that the map does not accurately display the exact boundaries of the search as defined in the paragraph above. The map is to be used for general reference purposes only.



A search of Heritage NSW AHIMS Web Services (Aboriginal Heritage Information Management System) has shown that:

3	Aboriginal sites are recorded in or near the above location.
0	Aboriginal places have been declared in or near the above location. *

If your search shows Aboriginal sites or places what should you do?

- You must do an extensive search if AHIMS has shown that there are Aboriginal sites or places recorded in the search area.
- If you are checking AHIMS as a part of your due diligence, refer to the next steps of the Due Diligence Code of practice.
- You can get further information about Aboriginal places by looking at the gazettal notice that declared it. Aboriginal places gazetted after 2001 are available on the [NSW Government Gazette \(https://www.legislation.nsw.gov.au/gazette\)](https://www.legislation.nsw.gov.au/gazette) website. Gazettal notices published prior to 2001 can be obtained from Heritage NSW upon request

Important information about your AHIMS search

- The information derived from the AHIMS search is only to be used for the purpose for which it was requested. It is not be made available to the public.
- AHIMS records information about Aboriginal sites that have been provided to Heritage NSW and Aboriginal places that have been declared by the Minister;
- Information recorded on AHIMS may vary in its accuracy and may not be up to date. Location details are recorded as grid references and it is important to note that there may be errors or omissions in these recordings,
- Some parts of New South Wales have not been investigated in detail and there may be fewer records of Aboriginal sites in those areas. These areas may contain Aboriginal sites which are not recorded on AHIMS.
- Aboriginal objects are protected under the National Parks and Wildlife Act 1974 even if they are not recorded as a site on AHIMS.
- This search can form part of your due diligence and remains valid for 12 months.